

60 Rec'd PCT/PRO-102 OCT 2000 OCT 2000

FORM PTO-1390 (REV 12-29-99)		U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE	ATTORNEY'S DOCKET NUMBER 43922
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371			U.S. APPLICATION NO. (if known, see 37 CFR 1.5) 09/623514
INTERNATIONAL APPLICATION NO. PCT/CA99/01202	INTERNATIONAL FILING DATE DECEMBER 16, 1999 ✓	PRIORITY DATE CLAIMED DECEMBER 17, 1978 ✓	
TITLE OF INVENTION DIACYLGLYCEROL ACYLTRANSFERASE GENE FROM PLANTS ✓			
APPLICANT(S) FOR DO/EO/US ZOU JITAO; TAYLOR, David C.; WEI, Yangdou; JAKO, Colette C.			
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:			
<ol style="list-style-type: none"> 1. <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. 371. 2. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371. 3. <input checked="" type="checkbox"/> This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1). 4. <input checked="" type="checkbox"/> A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date. 5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2)) <ol style="list-style-type: none"> a. <input type="checkbox"/> is transmitted herewith (required only if not transmitted by the International Bureau). b. <input checked="" type="checkbox"/> has been transmitted by the International Bureau. c. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US). 6. <input type="checkbox"/> A translation of the International Application into English (35 U.S.C. 371(c)(2)). 7. <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)) <ol style="list-style-type: none"> a. <input type="checkbox"/> are transmitted herewith (required only if not transmitted by the International Bureau). b. <input type="checkbox"/> have been transmitted by the International Bureau. c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired. d. <input checked="" type="checkbox"/> have not been made and will not be made. 8. <input type="checkbox"/> A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)). 9. <input checked="" type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). 10. <input type="checkbox"/> A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)). 			
Items 11. to 16. below concern document(s) or information included:			
<ol style="list-style-type: none"> 11. <input checked="" type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98. 12. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. 13. <input checked="" type="checkbox"/> A FIRST preliminary amendment. <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment. 14. <input type="checkbox"/> A substitute specification. 15. <input type="checkbox"/> A change of power of attorney and/or address letter. 16. <input checked="" type="checkbox"/> Other items or information: Sequence Listing on Diskette Statement of Conformity 			

Form PTO-1390 (REV 12-29-99) page 2 of 2

528 Rec'd PCT/PTO 02 OCT 20

Applicant :ZOU, Jitao et al.
Title : DIACYLGLYCEROL ACYLTRANSFERASE GENE FROM PLANTS

KIRBY EADES GALE BAKER,
Box 3432, Stn. D,
Ottawa, Ontario.
Canada K1P 6N9

The Hon. Commissioner of Patents
And Trademarks,
Washington DC 20231 U.S.A.

PRELIMINARY AMENDMENT

As a Preliminary Amendment, please amend this application as follows.

Page 1, immediately following line 1, add:

This application claims the priority rights of prior provisional U.S. Patent Application Serial No. 60/112,812 filed December 17, 1998, (now abandoned). --

The purpose of this amendment is merely to add a cross-reference to a related application.

Respectfully submitted,

John E.

Edwin Gale
Reg No. 28,584
Tel (613) 237-6900
Our File No. 43922
September 29, 2000

09/623514
528 Rec'd PCT/PTO 02 OCT 2000

File No.: 43922

IN THE UNITED STATES
PATENT AND TRADEMARK OFFICE

Applicant : ZOU, Jitao et al.
Title : DIACYLGLYCEROL ACYLTRANSFERASE GENE FROM
PLANTS
ATTENTION : BOX SEQUENCE

KIRBY EADES GALE BAKER,
Box 3432, Stn. D,
Ottawa, Ontario.
Canada K1P 6N9


The Hon. Commissioner of Patents
And Trademarks,
Washington DC 20231 U.S.A.

Dear Sir:

STATEMENT UNDER 37CFR § 1.825

I hereby state that the content of the paper copy of the Sequence Listing in the specification of the present application is the same as the content of the Sequence Listing in Computer-readable form on the diskette filed concurrently herewith.

Respectfully submitted,



Edwin Gale
Reg No. 28,584
Tel (613) 237-6900
Our File No. 42748
September 29, 2000

09/623514-1522960

13 Rec'd PCT/PTO 10 JUL 2001
09/623514

Attorney's Docket No. 43922

IN THE US PATENT AND TRADEMARK OFFICE

Applicant: ZOU, Jitao; et al.

Serial no.: 09/623,514

Filed: October 3, 2000

Title: DIACYLGLYCEROL ACYLTRANSFERASE GENE FROM PLANTS

Examiner:

Art Unit:

ATTENTION: BOX SEQUENCE

Kirby Eades Gale Baker
Box 3432 Stn. D
Ottawa, Ontario
Canada K1P 6N9

The Hon. Commissioner of Patents & Trademarks
Washington, DC 20231, USA

Sir:

Amendments to Sequence Listing and Computer Readable Copy thereof
under 37 C.F.R. § 1.825

This is with reference to the Notification of Missing Requirements mailed on May 14, 2001. Please amend this application as follows:

IN THE DISCLOSURE

Cancel the Sequence Listing (having its own page numbering scheme, pages 1 to 20), and replace this with the new Sequence Listing provided herewith as new pages 1 to 21.

A marked-up copy of the original Sequence Listing showing the amendments made at this time.

REMARKS

The Notice of Missing parts contained a Notification to Comply with Requirements for Patent Applications Containing Nucleotide Sequences and/or Amino Acid Sequence Disclosures.


A paper copy of an amended Sequence Listing and a copy in computer readable form on diskette are enclosed herewith.

This amendment corrects the Sequence Listing. This submission includes no new matter.

Also enclosed herewith is a statement of identity under 37 CFR § 1.821(f).

In addition, a copy of the Notice to Comply is attached.

Respectfully submitted



July 9, 2001

Edwin J. Gale
Reg. No. 28,584
Tel. (613) 237-6900
Fax (613) 237-0045
E-Mail: egale@kirbyeades.com

096254-0000

13 Rec'd PCT/PTO 10 JUL 2001

Attorney Docket No. 43922
09/623514

IN THE US PATENT AND TRADEMARK OFFICE

Applicant: ZOU, Jitao; et al.

Serial no.: 09/623,514

Filed: October 3, 2000

Title: DIACYLGLYCEROL ACYLTRANSFERASE GENE FROM PLANTS

Examiner:

Art Unit:

ATTENTION: BOX SEQUENCE

Kirby Eades Gale Baker
Box 3432 Stn. D
Ottawa, Ontario
Canada K1P 6N9

The Hon. Commissioner of Patents & Trademarks
Washington, DC 20231, USA

Sir:

Statement under 37 CFR § 1.821(f)

This is with reference to the accompanying response of even date to the Notice to Comply mailed on May 14, 2001.

The undersigned hereby states that the copy of the Sequence Listing in computer readable form on the attached diskette is the same as the substitute copy of the Sequence Listing enclosed in the aforesaid response.

July 9, 2001

Edwin J. Gale
Reg. No. 28,584
Tel. (613) 237-6900
Fax (613) 237-0045
E-Mail: egale@kirbyeades.com

Respectfully submitted,



DIACYLGLYCEROL ACYLTRANSFERASE GENE FROM PLANTSTECHNICAL FIELD

This invention relates to plant genes useful for the genetic manipulation of plant characteristics. More specifically, the invention relates to the identification, 5 isolation and introduction of diacylglycerol acyltransferase (DGAT) genes useful, for example, for altering the seed oil content, the ratio of diacylglycerol / triacylglycerol proportions in the seed oil, fatty acid synthesis, seed oil acyl composition, seed size/weight and carbon flux into other seed components, in commercial or crop plants.

10 BACKGROUND ART

Plant seed oils are major sources of essential polyunsaturated fatty acids for human diets and renewable feedstocks for chemical industries. The enzymes of the fatty acid synthase complex in the plastids of developing seeds are responsible for the biosynthesis of fatty acids that are channeled into the cytosolic acyl-CoA pool to 15 sustain triacylglycerol accumulation. Triacylglycerol (TAG) biosynthesis is located in the endoplasmic reticulum with glycerol-3-phosphate and fatty acyl-CoAs as the primary substrates. There are three acyltransferases involved in the plant storage lipid bioassembly, namely the glycerol-3-phosphate acyltransferase (GPAT, EC 2.3.1.15), the lyso-phosphatidic acid acyltransferase (LPAT, EC 2.3.1.51) and the 20 diacylglycerol acyltransferase (DGAT, EC 2.3.1.20). These three acyltransferases catalyze the stepwise acylation of the glycerol backbone with the final step being the acylation of sn-1, 2-diacylglycerol (DAG) by DGAT into the formation of TAGs, a biochemical process generally known as the Kennedy pathway (Stymne and Stobart, 1987).

25 Among the three ER-based fatty acyl-CoA acyltransferases, only LPAT gene(s) have been cloned from plants (Knutzon *et al.*, 1995, Lassner *et al.*, 1995). Like several other enzymes involved in storage lipid biosynthesis, acyltransferases are intrinsic ER proteins and are extremely difficult to purify. The research on plant DGAT has been largely limited to studies of activity profiles by using the particulate 30 fractions generated by differential centrifugation of seed or microspore-derived embryo homogenates (Weselake *et al.*, 1993). Although partial purification of

FO6260-4T5E2960

DGAT from cotyledons of germinating soybean seeds was reported (Kwanyan and Wilson, 1986), detailed molecular characterization of this enzyme is lacking.

Reference is made to R61u012 Database Entry Ac005917; Accession number AC005917; 4 November 1998; LIN X. ET AL : "Arabidopsis thaliana chromosome II section 113 of 255 of the complete sequence". This relates to a nucleotide sequence deposit first submitted to the NCBI GenBank on November 3, 1998 without any identification of putative coding sequences contained therein. There was no reference in the deposited materials to a "diacylglycerol O-acyltransferase gene".

Reference is also made to VESNA KATAVIC ET AL.: "Alteration of Seed Fatty Acid Composition by an Ethyl Methanesulfonate-Induced Mutation in Arabidopsis thaliana Affecting Diacylglycerol Acyltransferase Activity"; PLANT PHYSIOLOGY, vol. 108, 1995, pp. 399-409. This reference discloses an Arabidopsis mutant designated AS11 that has reduced diacylglycerol acyltransferase activity. The reference does not disclose any DNA sequences whatsoever and teaches only that alterations in DGAT activity may lead to changes in fatty acid content.

Accordingly, while the Kennedy pathway is known and shows the steps in the biosynthesis of TAGs in plants, there has not been any identification and use of a genetic element that can be used reliably in plants to modify TAG synthesis and composition in a way that may be exploited commercially.

DISCLOSURE OF THE INVENTION


An object of the invention is to identify, isolate and clone a genetic element that may be used to modify the natural formation of triacylglycerols in plants in order to increase the yield of commercial plant oils, or to modify their composition to achieve specific commercial improvements of plants and plant products.

Another object of the invention is to identify, isolate and characterize diacylglycerol acyltransferase (DGAT) gene and cDNA sequences from Arabidopsis and to utilize these sequences in the genetic manipulation of plants.

Another object of the invention is to provide a vector containing the full-length DGAT cDNA sequence from *Arabidopsis* in a sense orientation under the control of a seed-specific promoter (e.g. napin; See Josefsson et al., 1987; Radke et al., 1988; Voelker et al., 1992), for re-introducing into *Arabidopsis* or for
5 introducing into other plants.

Another object of the invention is to provide a vector containing a genomic fragment from *Arabidopsis* consisting of the full-length DGAT gene under the control of its own 5' upstream regulatory sequences, for re-introducing into *Arabidopsis* or for introducing into other plants.

10 Another object of the invention is to provide a method to construct a vector containing the full-length DGAT sequence or a significant portion of the DGAT sequence from *Arabidopsis*, in an antisense orientation under control of either a constitutive or a seed-specific promoter, for re-introducing into *Arabidopsis* or for introducing into other plants.



AMENDED SHEET

Another object of the invention is to provide a method of modifying *Arabidopsis* and other plants to change their seed oil content.

Another object of the invention is to provide a method of modifying *Arabidopsis* and other plants to change the acyl composition of their seed oil.

- 5 Another object of the invention is to provide a method of modifying *Arabidopsis* and other plants to change their average seed weight or size.

According to one aspect of the present invention, there is provided a vector containing isolated and purified deoxyribonucleic acid (cDNA) of SEQ ID NO:1 (pDGATcDNA; ATCC No PTA-989), for introduction of the cDNA in a sense
10 orientation into a plant cell.

As another aspect of the present invention, there is provided a vector containing isolated and purified genomic deoxyribonucleic acid (genomic DNA) of SEQ ID NO:3 (pDGATgene; ATCC No PTA-988), for introduction of the gene in a sense orientation into a plant cell.

- 15 According to yet another object of the invention, there is provided a method for preparing a vector containing SEQ ID NO:1 or a part thereof, for introduction of the gene or partial gene in an antisense orientation, into a plant cell.

According to yet another object of the invention, there is provided seed of *Arabidopsis thaliana* ecotype Columbia mutant AS11 (ATCC No. PTA-1013) and
20 characterization of its lipid phenotype (Katavic et al., 1995; Zou et al. 1999). The AS11 mutant seed line has an insertion mutation at the *TAG1* locus on chromosome II, and produces plants exhibiting reduced DGAT activity (Figure 4) and an reduced TAG/DAG ratio during seed development (Table 1), resulting in an altered seed fatty acyl composition (Figure 2), reduced oil content (Table 1), and
25 increased seed oil diacylglycerol content during development (Figure 3) and at maturity (lower TAG/DAG ratio of Table 1). The cDNA sequence of the AS11 DGAT is shown in SEQ ID NO:23, the genomic DNA sequence is shown in SEQ ID NO:24 and the translated protein sequence of the AS11 DGAT is shown in SEQ ID NO:25.

The invention also relates to transgenic plants and plant seeds having a genome containing an introduced DNA sequence of SEQ ID NO:1 or SEQ ID NO:3, and a method of producing such plants and plant seeds.

The invention also relates to SEQ ID NO:1 or SEQ ID NO:3, or a part of
5 SEQ ID NO:1 or SEQ ID NO:3, or SEQ ID NO:1 containing an 81 bp insertion [SEQ ID NO: 23] or SEQ ID NO:3 containing an 147 bp insertion [SEQ ID NO:24] such that the deduced amino acid sequence of the encoded protein contains the repeated sequence SHAGLFNLCVVVLIAVNSRLIIENLMK [SEQ ID NO:25] where the spacing and identity of the underlined amino acids are identical or are replaced
10 by conserved substitutions, characterized in that said sequence has been introduced in sense or antisense orientation, and a method of producing such plants and plant seeds.

As will be appreciated by persons skilled in the art, the invention also relates to substantially homologous DNA sequences from plants encoding proteins with
15 deduced amino acid sequences of 25% or greater identity, and 40% or greater similarity, isolated and/or characterized and/or designed by known methods using the sequence information of SEQ ID NO:1 or SEQ ID NO:3 or or SEQ ID NO:1 containing an 81 bp insertion [SEQ ID NO:23] such that the deduced amino acid sequence of the encoded protein contains the repeated sequence
20 SHAGLFNLCVVVLIAVNSRLIIENLMK (SEQ ID NO:25) where the spacing and identity of the underlined amino acids are identical or are replaced by conserved substitutions, and to parts of reduced length that are still able to function as inhibitors of gene expression by use in an anti-sense, co-suppression (Transwitch; Jorgensen and Napoli 1994) or other gene silencing technologies. It will be
25 appreciated by persons skilled in the art that small changes in the identities of nucleotides in a specific gene sequence may result in reduced or enhanced effectiveness of the genes and that, in some applications (e.g. anti-sense or co-suppression), partial sequences often work as effectively as full length versions. The ways in which the gene sequence can be varied or shortened are well known to
30 persons skilled in the art, as are ways of testing the effectiveness of the altered genes. All such variations of the genes are therefore claimed as part of the present invention.

Other preferred degrees of identity to the indicated sequences are at least 30%, 40%, 50%, 60%, 70%, 80%, 90% and 95%; and other preferred degrees of similarity are at least 50%, 60%, 70%, 80%, 90% and 95%. The inventors have used a computer program known as MegAlign®, DNASTAR® of DNASTAR Inc.,
5 1228 South Park Street, Madison, WI 53715, USA, for assessing homology. This program is based on the Clustal V algorithm (Higgins and Sharp (1998): A package for performing multiple sequence alignment on a microcomputer; GENE 73:237-244). For each gap introduced in the alignment, the program deducts a penalty from the score. A higher gap penalty suppresses gapping; a lower value promotes
10 it. The program also assesses penalties based on the length of the gap. The more residues the gap spans, the greater the penalty. The program deducts these penalties from the overall score of the alignment.

Stated more generally, the present invention relates to the isolation, purification and characterization of a diacylglycerol acyltransferase (DGAT) gene
15 from the *Brassicaceae* (specifically *Arabidopsis thaliana*) and demonstrates its utility in regulating fatty acid synthesis, seed oil content, diacylglycerol/triacylglycerol ratios and seed size/weight. Until now, no concrete data is available on the gene structure of plant DGATs, or their utility in altering oil content or composition through genetic manipulation.

20 When considering altered oil contents or compositions, results from averages of statistically-significant numbers of plants or seeds according to the invention are best compared with results from averages of statistically-significant numbers of untransformed (control) plants or seeds of the same genotype grown under identical conditions at the same time. This allows for the variability of
25 individual plants of the same genotype, particularly when such plants are grown under different conditions. The actual number of plants or seeds used to form the required average may vary, but should be enough to provide a generally constant average whenever such number is selected. Generally, the number should be at least 10, and is more preferably at least 20, 30, 50 or 100.

30 The DGAT gene was cloned, characterized and authenticated from *Arabidopsis* by: (1) selection and characterization of plant ESTs sharing some homology to mammalian acyl-CoA: cholesterol acyltransferases; (2) the functional

expression of a full-length cDNA in yeast; (3) the characterization and isolation of the DGAT (*TAG1*) gene from *Arabidopsis* mutant AS11 containing an insertion mutation in the DGAT gene and a seed oil phenotype which consists of an altered DAG/TAG ratio, and an altered oil content and acyl composition; (4) complementation of the AS11 mutant by
 5 Insertion of the DGAT cDNA sequence to restore a wild-type fatty acid composition; (5) the over-expression of the DGAT cDNA in wild-type *A. thaliana* transgenic plants which produce seeds with an increased oil content, increased average seed weight and altered seed oil acyl composition.

The *A. thaliana* DGAT structure is significantly homologous (over 40% identical
 10 over a region of more than 400 amino acids) to its mammalian counterparts, and is highly homologous to subsequently reported putative *B. napus* DGATs at both the nucleotide (92%) and the deduced amino acid (90%) levels (Nykiforuk et al, 1999; GenBank/EMBL Accession No AF155224; AF164434).

The DGAT of the current invention is useful in manipulating DGAT activity, and triacylglycerol bioassembly in plants. For example, by transforming plants with a construct
 15 containing the DGAT gene in a sense orientation, under the control of a tissue-specific promoter (e.g. seed-specific promoter napin), the expression of DGAT and accumulation of seed oil can be enhanced or the acyl composition of the seed oil altered. Yet another example would be to express the DGAT cDNA under the control of a constitutive promoter
 20 (e.g. 35S (Datla et al., 1993)) to increase the TAG content of vegetative tissues (leaves, roots, stems). This may have particular advantages for altering the starch/oil ratio in root crops.

Alternatively, DGAT expression can be silenced to some degree by anti-sense or co-suppression (Transwitch) phenomena (De Lange et al., 1995; Mol et al., 1990;
 25 Jorgensen and Napoli, 1994; Kinney, 1995; Vaucheret et al, 1998; Taylor, 1998). For example, silencing DGAT in a seed specific manner may result in a reduction in TAG accumulation. This could have applications in reducing the oil content in seed barley to enhance stability during storage. As a second example, seed-specific silencing may lead to a relatively high accumulation of DAG or an increase in the DAG/TAG ratio in the
 30 developing or mature seed. As yet another example, the expression of the mutated DGAT gene which results in a 27 amino

acid repeat insertion in the mutant DGAT protein (See Figure 5 a) can be used to alter the DAG/TAG ratio in developing and mature seed. Such manipulations can lead to edible seed oils produced naturally in the plant, containing enhanced relative levels of DAG /reduced levels of TAG (See Figure 3; Table 1) to act as all-natural

5 emulsifiers in the food and confections industries, or to enhance the nutritional/health profile of vegetable oils as functional foods (e.g. as cooking oils, stir fry oils, in salad dressings, margarines etc.) by inhibiting neutral fat deposition in humans. Processed oils produced from canola and soybean which contain increased proportions of diacylglycerol have been cited by the Kao Corporation of

10 Japan (e.g. Econa Cooking Oil; Kao Corporation KI, 1-14-10 Nihonbashi-Kayabacho, Chuoku, Tokyo 103 Japan; e-mail: 210064@kastanet.kao.co.jp) as a product making it difficult for blood neutral fat to increase after a meal, and for fat to cling to the body, thereby assisting individuals who are overweight or who suffer from high neutral fat levels. As a third example, silencing or reducing the activity of

15 DGAT in a seed specific manner (as observed in mutant AS11; e.g. by over-expressing SEQ ID NO:23 or silencing expression of SEQ ID NO:1 or SEQ ID NO:3), and combining this trait with the capacity to produce polyhydroxyalkanoates (PHAs; e.g. polyhydroxybutyrate) in seeds (Poirier et al., 1992; 1995) will allow an increased flow of unesterified fatty acids towards β -oxidation (Poirier et al., 1999).

20 By recycling or diverting the unesterified fatty acids into β oxidation, the resulting acetyl-CoA moieties will lead to a significant increase in polyhydroxyalkanoates (PHAs) or a change in PHA composition (Poirier et al., 1999). Transgenic plants producing PHAs in seeds have potential for utility as biodegradable thermoplastics. However, up to now, the levels of PHAs produced have been relatively small

25 (Poirier et al, 1992; 1995). The utility of transgenically reducing the DGAT activity to significantly enhance PHA production (e.g. 10-fold increase) in PHA-producing seeds is now possible, due to the current DGAT invention.

Some of the manipulations and deliverables which are possible using the DGAT gene or a part thereof, include, but are not limited to, the following: seeds with

30 increased or decreased oil content; seeds containing oils with an enhanced diacylglycerol content, seed oils with an altered acyl composition; plants producing larger or heavier seeds; plants exhibiting an enhanced or altered capacity to accumulate storage compounds in other storage organs (e.g. tubers, roots).

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a diagram illustrating the Kennedy pathway for the bioassembly of triacylglycerols in plants, and shows the critical role played by DGAT as the final step of the Kennedy pathway.

5 Figure 2 is a graph showing the comparison of the fatty acid composition of seed oil from wild-type (WT) and DGAT mutant AS11 lines of *Arabidopsis thaliana*. Proportions of fatty acids are reported as Mol % of the total fatty acid composition of the seed oil from each line. Error bars are \pm SE (n= 10 plants of each line sampled, with 50 seeds per sample per analysis).

10 Figure 3 is a graph showing a comparison of the DAG content in developing seed of wild-type (WT) and DGAT mutant AS11 lines of *Arabidopsis thaliana*. More specifically, it is a comparison of the fatty acid content of DAG pool in wild-type green developing seeds compared to that of the AS11 mutant.

Figure 4 is a graph showing a comparison of the DGAT activity
15 (pmol/min/mg protein) in developing seeds at the milky, early green, mid-green and brown-green stages of embryo development in wild-type (WT) and DGAT mutant AS11 lines of *Arabidopsis thaliana*. Developing seeds at each stage were selected and DGAT enzyme analyses conducted as described previously by Katavic et al., 1995).

20 Figure 5(a) shows the amino acid sequence alignment of the *Arabidopsis* DGAT (AtTAG1) [SEQ ID NO:2] with mammalian (mouse and putative human) DGATs. MDGAT, mouse DGAT [SEQ ID NO:4]; GenBank/EMBL Accession No. AF078752 (Cases et al., 1998); HARGP1, human ARGPI protein [SEQ ID NO:5]; GenBank/EMBL Accession No. AF059202; Oelkers et al., 1998]. Dots indicate
25 gaps. Identical amino acid residues are highlighted in black. Conserved residues are shaded. The 27-amino acid repeat found in *Arabidopsis thaliana* mutant AS11 and generated by the insertion mutation (81 bp) found in SEQ ID NO:23 (SHAGLFNLCVVVLIIVNSRLIENLMK) is overlined thus: ----- . The putative diacylglycerol binding site is overlined thus: =====. The SnRK1 targeting site is
30 overlined thus: ===== with an asterisk (*) over the serine (S) residue as the targeted phosphorylation site.

Figure 5(b) shows the Kyte-Doolittle hydropathy plot of the DGAT protein.

Figure 6(a) shows the results of a Northern analysis of *TAG1* gene expression in *Arabidopsis thaliana*. Total RNA was extracted from roots (RT), leaves (LF), flowers (FL), young seedlings (YS), developing siliques (SL), and germinating seeds (GS).

Figure 6(b) shows the results of Southern blot analysis of the *TAG1* gene in *Arabidopsis thaliana*. Genomic DNA was digested with restriction enzymes BglII (Lane 1), EcoRI (Lane 2), and HindIII (Lane 3). The *TAG1* DNA probe was ³²P labeled by random priming.

Figure 7(a) is a diagrammatic representation of the *TAG1* gene structure. The boxes indicate the 16 exons (solid boxes for coding regions, open box for untranslated regions), and the lines represent the 15 introns. A, B and C denote the positions of the primers used for PCR amplifications of the segments from wild type (WT) and AS11. The specific primers A, B and C are described in Experimental Procedures: *Primer Strategy* (found later in this specification).

Fig 7(b) shows gel separation of the PCR products amplified from wild type (WT) and AS11. Lane 1, PCR product with primers A and B using WT genomic DNA as template. Lane 2, PCR product with primers A and B using AS11 genomic DNA as template. Lane 3, PCR product with primers C and B using WT genomic DNA as template. Lane 4, PCR product with primers C and B using AS11 genomic DNA as template. Lane 5, RT-PCR with primers A and B using RNA prepared from WT seedling RNA. Lane 6, RT-PCR with primers A and B using RNA prepared from AS11 seedling RNA.

Figure 8 is a graph showing microsomal DGAT activity in Yeast Strain YMN5 Transformed with empty plasmid (pYES Con) and with the *A. thaliana* DGAT cDNA (pYES:DGAT). This illustrates the expression of the *TAG1* cDNA in yeast. Host cultures of strain YMN5 were transformed with pYES2 plasmid only (pYES2; without *TAG1* insert) or with pYES2 containing the *TAG1* cDNA insert (pYES2:*TAG1*). Following induction in the presence of galactose, transformants were lysed and assayed for DGAT activity as described in the Experimental Procedures. The results of two separate DGAT activity experiments are shown.

Figure 9 is a map of plasmid pSE129A which may be used as a vector. The vector contains the following salient features for plant transformation in the current invention: the seed-specific napin promoter and NOS terminator between which is a multiple cloning site.

5 Figure 10 is a graph showing the complementation of the AS11 DGAT mutation with the wild-type cDNA. Transformation of *Arabidopsis thaliana* mutant line AS11 with the DGAT cDNA [SEQ ID NO:1] under the control of a napin promoter, leads to a restoration of the wild-type (WT) fatty acid composition in the seed oil of the transformant lines 3-4, 9-1, 14-2 and 9-4. Fatty acid composition (wt
10 %) was determined on the seed oil extracted from 100-seed samples from *A. thaliana* non-transformed controls (WT), mutant line AS11, and T₂ seeds of napin:DGAT transgenic lines.

Figure 11 is a graph showing the seed oil content of non-transformed WT control, and pRD400 control (empty plasmid) and napin:DGAT T₂ transgenic
15 *Arabidopsis thaliana* seed lines. More particularly, the graph shows the transformation of wild type (WT) *Arabidopsis thaliana* with the DGAT cDNA [SEQ ID NO:1] under the control of a napin promoter, leads to a higher seed oil content in the DGAT transgenic lines. Oil content is expressed as µg total fatty acids (FAs) per
20 100 seeds from *A. thaliana* non-transformed controls (WT Con), and T₂ seeds of pRD400 control (empty plasmid) transgenic, and napin:DGAT transgenic lines 1', 2', 9, 10 and 11. Standard error bars for the control lines are indicated; n=10.

Figure 12 is a graph showing the average 100-seed weight of non-transformed WT control, and pRD400 control (empty plasmid) and napin:DGAT T₂ transgenic *Arabidopsis thaliana* seed lines. More specifically, the graph shows the
25 over-expression of the DGAT cDNA under the control of a napin promoter, in wild-type (WT) *Arabidopsis thaliana* leads to a higher average seed weight. The average weight of 100-seed samples from *A. thaliana* non-transformed controls (WT Con), and T₂ seeds of pRD400 control (empty plasmid) transgenic, and napin:DGAT transgenic lines 1', 2', 9, 10 and 11 are reported as mg dry weight (DW).

30 Figure 13 is a graph showing the positive correlation between oil content (expressed as µg Total fatty acids (FAs) per 100 seeds) and average seed weight (expressed as average mg DW per 100-seed samples) from *A. thaliana* non-

transformed controls (WT Con ♦), and T2 seeds of pRD400 control (empty plasmid) transgenic (○) and napin:DGAT transgenic lines 1', 2', 9, 10 and 11 (■) are reported as mg dry weight (DW).

BEST MODES FOR CARRYING OUT THE INVENTION

- 5 Fig. 1 is a diagram illustrating the Kennedy pathway for the biosynthesis of TAGs in plants. Of the various illustrated enzymes, DGAT is the only enzyme in the Kennedy pathway that is exclusively committed to TAG biosynthesis, and its key role is apparent from the scheme of Figure 1. *sn*-1,2-DAG, generated as a result of either the catalytic action of PA phosphatase (EC 3.1.3.4) or CPTase (EC 2.7.8.2),
10 can be used in the biosynthesis of TAG.

For this reason, the inventors of the present invention decided to investigate DGAT to see if the corresponding gene in plants could be sequenced and cloned and used to modify the seed oil content and fatty acid composition of plants in a way that could be commercially useful.

- 15 The acyl-CoA dependent acylation of *sn*-1,2-DAG is catalyzed by DGAT (Stymne and Stobart, 1987). In developing and germinating seeds of oilseed plants, TAG accumulation and DGAT activity have been shown to associate with the endoplasmic reticulum (ER; high speed microsomal fraction) (Stobart et al., 1986; Cao and Huang, 1986; Stymne and Stobart, 1987; Frentzen, 1993; Settlage et al,
20 1995; Lacey and Hills, 1996). The biochemical properties of microsomal DGAT have been examined in a number of plant systems (Frentzen, 1993) including developing seeds (Bernerth and Frentzen, 1990; Vogel and Browse, 1996; Cao and Huang, 1987) and embryo cultures (Taylor et al., 1991; Weselake et al., 1991; Taylor et al., 1992; Little et al., 1994) of *B. napus* L. In general, studies with developing seeds
25 indicate that DGAT activity increased rapidly during the active phase of oil accumulation and then decreases markedly as seed lipid content reaches a plateau (Tzen et al., 1993; Weselake et al., 1993).

- A number of studies with both mammalian (Mayorek et al., 1989; Tijburg et al., 1989) and plant (Ichihara et al., 1988; Perry and Harwood, 1993 a and 1993 b;
30 Settlage et al. 1995) systems have suggested that DGAT may catalyze a rate-limiting reaction in TAG bioassembly. However, this hypothesis has not been

rigorously tested, and has not been reduced to practice by transgenic expression of any DGAT gene in plant or animal systems, until now. Developing seeds of *B. napus* L., cv Shiralee, have been shown to produce significant levels of DAG during the active phase of oil accumulation suggesting that DGAT catalyzed

5 reaction may regulate the flow of carbon into TAG (Perry and Harwood, 1993 a and 1993 b). In addition, an ethyl methanesulfonate-induced (EMS-induced) mutant of *A. thaliana*, designated AS11, has been shown to have a reduced DGAT activity that correlated with both an increased DAG pool and decreased accumulation of TAG (Katavic et al. 1995). Given its possible rate-limiting role in TAG bioassembly,

10 the inventors of the present invention have identified DGAT as a potential target in the genetic modification of plant lipid biosynthesis.

Previously, the partial characterization of an EMS-induced *Arabidopsis thaliana* mutant, AS11, with altered fatty acid composition was reported (Katavic et al., 1995). In comparison to wild type plant seeds, AS11 seeds have reduced levels

15 of the very long chain fatty acid eicosenoic acid (20:1) and reduced oleic acid (18:1) and accumulate α -linolenic acid (18:3) as the major fatty acid in triacylglycerols (Figure 2). The AS11 mutant has a consistently lower ratio of TAG/DAG in developing seeds, and it accumulates an elevated amount of seed DAG (Figure 3), the substrate of the diacylglycerol acyltransferase. Through a series of biochemical

20 analyses, it was shown that AS11 had reduced diacylglycerol acyltransferase activities throughout seed development (Figure 4). AS11 also had a reduced (by 25-30%) oil content phenotype, providing some evidence that DGAT may be controlling flux into TAG biosynthesis, as shown in Table 1 below. The AS11 did not have a wrinkled-seed phenotype as described in other low-seed-oil mutants (Focks

25 and Benning, 1998).

Table 1

Comparison of AS11 [Katavic et al., (1995) and wild-type *A. thaliana* seeds with respect to lipid profiles at mid-development, and the relative TAG, DAG and sterol ester contents in AS11 and WT seeds at maturity.

Seed Type	TAG/DAG ratio at mid- development ^a	TAG/DAG ratio at maturity ^a	Relative TAG content at maturity ^b °(nmol/mg DW)	Sterol Esters at maturity (% of Total Lipid Extract) ^d
WT	17	90	1.00 ^b °(255)	0.8
AS11	5	20	0.6 ^b °(174)	1.15

^a Embryos staged and lipids measured as described in Katavic et al., (1995);

^b Relative TAG content of 200-seed samples of AS11 and WT were measured by MASS-1H-NMR according to the method of Rutar (1989). The integration response for resonances attributable to liquid-like oil were summed and the value for AS11 seed is reported relative to the response for the WT control seed sample (the latter set at a value of 1.00); ^c TAG content (nmoles/mg DW) measured by transmethylation of a TLC-purified TAG fraction, followed by GC analysis of fatty acid methyl esters; ^d A total lipid extract was prepared as described by Taylor et al., (1991; 1992), and sterol esters isolated and characterized as described in the Experimental Procedures.

Genetic analysis indicated that the fatty acid phenotype is caused by a semidominant mutation in a nuclear gene, designated *TAG1*. The mutation was mapped to chromosome II, and was estimated to lie in the region approximately 17.5 ± 3 cM from the *sti* locus and 8 ± 2 cM from the *cp2* locus.

- 5 Because a DGAT gene has not heretofore been cloned from any plant, until now, it has not been possible to address the possibility of genetic modifications to alter carbon flux, increase fatty acid synthesis, oil content, oil acyl composition, or seed size, by modulating plant DGAT activity.

- 10 However, there are many examples of successful modifications to plant metabolism that have been achieved by genetic engineering to transfer new genes or to alter the expression of existing genes, in plants. It is now routinely possible to introduce genes into many plant species of agronomic significance to improve crop performance (e.g. seed oil or tuber starch content/composition; meal improvement;

herbicide, disease or insect resistance; heavy metal tolerance etc.) (MacKenzie and Jain, 1997; Budziszewski et al., 1996; Somerville, 1993; Kishore and Somerville, 1993).

For example, increases in the proportions of some strategic fatty acids and
5 in the quantities of seed oil have been achieved by the introduction of various fatty acid biosynthesis and acyltransferase genes in oilseed crops. These include the following demonstrations: Expression of an anti-sense construct to the stearoyl-ACP $\Delta 9$ desaturase in Brassicaceae led to an increase in the stearic acid content (Knutzon et al., 1992). Expression of a medium chain fatty acyl-ACP thioesterase
10 from California Bay, in Brassicaceae was demonstrated to increase the lauric acid (12:0) content (Voelker et al., 1992; 1996). Expression of a Jojoba β keto-acyl-CoA synthase in low erucic acid Brassicaceae led to an increase the level of erucic acid (22:1); the effect following expression in high erucic acid cultivars was negligible (Lassner et al., 1996). Increased proportions of oleic acid in Brassica napus and in
15 soybean have been achieved by silencing the microsomal FAD2 ($\Delta 12$) desaturase (Hitz et al., 1995; Kinney, 1995 ; 1997). Transformation of Arabidopsis thaliana and rapeseed (B. napus) with a yeast sn-2 acyltransferase resulted in seed oils with increased proportions of 22:1 and other very long-chain fatty acids and significant increases in seed oil content (Zou et al., 1997).

20 Starch deposition has also been altered by genetic engineering. By expression of a mutant E. coli glgC16 gene encoding an ADP glucose pyrophosphorylase in potato tubers, an increase in starch accumulation was achieved (Stark et al., 1992).

The inventors therefore considered the DGAT gene to hold great promise for
25 the desired modification of TAGs in plants.

The best modes for carrying out the invention will be apparent from the following description of the results of tests and experiments that have been carried out by the inventors.

The inventors chose to use the well-accepted model plant system
30 *Arabidopsis thaliana* for the cloning of DGAT, as a host system for genetic engineering to alter DGAT expression, and to study the effects of altering DGAT

- expression on seed triacylglycerol bioassembly. This is because, over the past several years, *Arabidopsis thaliana*, a typical flowering plant, has gained increasing popularity as a model system for the study of plant biology. As a result of the ease with which this plant lends itself to work in both classical and molecular genetics,
- 5 *Arabidopsis* has come to be widely used as a model organism in plant molecular genetics, development, physiology and biochemistry (Meyerowitz and Chang, 1985; Meyerowitz, 1987; Goodman et al., 1995). This model dicotyledonous plant is also closely related to *Brassica* crop species and it is increasingly apparent that information concerning the genetic control of basic biological processes in
- 10 *Arabidopsis* will be transferable to other species (Lagercrantz et al., 1996).

- Indeed, there are numerous examples wherein studies of the molecular biology and biochemistry of a particular metabolic pathway or developmental process and the possibility of genetically engineering a plant to bring about changes to said metabolic pathway or process, has first been tested in the model plant
- 15 *Arabidopsis*, and then shown to yield similar phenotypes in other plants, particularly crop plants.

- For example, the extra- plastidial membrane associated oleate (18:1) $\Delta 12$ (ω -6) desaturase gene, *FAD2*, was originally studied and eventually cloned from *Arabidopsis thaliana*, by identifying the lesion found in an *A. thaliana* mutant
- 20 defective in desaturating oleate to produce linoleate (18:2) on the phosphatidylcholine backbone. This resulted in a high oleic acid phenotype in the *A. thaliana* seed oil (Okuley et al., 1994). Genetic engineering of both soybean (*Glycine max.*) and canola *B. napus* to silence the indigenous *FAD2* gene(s) in a seed-specific manner by anti-sense or co-suppression approaches, resulted in
- 25 similar high oleic acid seed oil phenotypes (Kinney, 1995; 1997).

- Transgenic expression of a yeast *sn-2* acyltransferase (*SLC1-1*) gene to achieve enhanced seed oil and very long-chain fatty acid content was first performed in *Arabidopsis* and later shown to yield similar phenotypes in transgenic rapeseed (*B. napus*) experiments (Zou et al., 1997). *Arabidopsis thaliana* has
- 30 repeatedly shown itself to be a useful model system for metabolic engineering of metabolic pathways (e.g. lipid biosynthesis, photosynthesis) or processes (organogenesis, reproductive development etc.) common to all higher plants.

In the area of secondary metabolism/signal transduction, an anthocyanin pathway-specific transcriptional activator from the monocot maize designated as *R* (the *myc* transcription factor involved in activation of biosynthetic genes for anthocyanin production in the aleurone cells of maize kernels), was expressed in
5 the dicot *Arabidopsis*, causing augmented anthocyanin pigmentation in the inflorescences. Subsequent expression in another dicot, tobacco (*Nicotiana tabacum*), resulted in similar floral pigmentation changes (Lloyd et al., 1992). These experiments demonstrate that whole pathways common to all flowering plants can be co-ordinately controlled through the introduction of transcriptional regulators, and
10 that the mechanisms are common to diverse plant species.

In the context of the current invention, all plant seeds accumulate some triacylglycerol (oil) and this ubiquitous process is affected, at least in part, by the activity of a microsomal DGAT, as explained previously. Thus, many of the effects observed following genetic engineering to modulate DGAT expression in
15 *Arabidopsis* can be expected and predicted to result in similar phenotypes when carried out in all other plants. For example, after the present invention was made, information has become available that supports the findings of the present inventors by showing that *B. napus* has a highly homologous DGAT gene (Nikiforuk et al., 1999), and thus *B. napus* is a clear target for similar genetic modifications as those
20 shown for *A. thaliana*.

There are a number of ways by which genes and gene constructs can be introduced into plants, and a combination of plant transformation and tissue culture techniques have been successfully integrated into effective strategies for creating transgenic crop plants. These methods, which can be used in the present
25 invention, have been extensively reviewed elsewhere (Potrykus, 1991; Vasil, 1994; Walden and Wingender, 1995; Songstad et al., 1995), and are well known to persons skilled in the art. For example, one skilled in the art will certainly be aware that, in addition to *Agrobacterium*-mediated transformation of *Arabidopsis* by vacuum infiltration (Bechtold et al., 1993) or wound inoculation (Katavic et al.,
30 1994), it is equally possible to transform other plant and crop species, using *Agrobacterium* Ti-plasmid-mediated transformation (e.g. hypocotyl (DeBlock et al., 1989) or cotyledonary petiole (Moloney et al, 1989) wound infection), particle bombardment/biolistic methods (Sanford et al., 1987; Nehra et al., 1994; Becker et

al., 1994) or polyethylene glycol-assisted protoplast transformation (Rhodes et al., 1988; Shimamoto et al., 1989) methods.

As will also be apparent to persons skilled in the art, and as extensively reviewed elsewhere (Meyer, 1995; Datla et al., 1997), it is possible to utilize plant promoters to direct any intended up- or down-regulation of transgene expression using constitutive promoters (e.g. those based on CaMV35S), or by using promoters which can target gene expression to particular cells, tissues (e.g. napin promoter for expression of transgenes in developing seed cotyledons), organs (e.g. roots), to a particular developmental stage, or in response to a particular external stimulus (e.g. heat shock).

Particularly preferred plants for modification according to the present invention include *Arabidopsis thaliana*, borage (*Borago* spp.), Canola, castor (*Ricinus communis*), cocoa bean (*Theobroma cacao*), corn (*Zea mays*), cotton (*Gossypium* spp.), *Crambe* spp., *Cuphea* spp., flax (*Linum* spp.), *Lesquerella* and *Limnanthes* spp., Linola, nasturtium (*Tropaeolum* spp.), *Oenothera* spp., olive (*Olea* spp.), palm (*Elaeis* spp.), peanut (*Arachis* spp.), rapeseed, safflower (*Carthamus* spp.), soybean (*Glycine* and *Soja* spp.), sunflower (*Helianthus* spp.), tobacco (*Nicotiana* spp.), *Vernonia* spp., wheat (*Triticum* spp.), barley (*Hordeum* spp.), rice (*Oryza* spp.), oat (*Avena* spp.) sorghum (*Sorghum* spp.), rye (*Secale* spp.) or other members of the *Gramineae*.

The present invention is particularly useful when used to modify the yield or composition of oilseed produced from oilseed crops. Oilseed crops are plant species that are capable of generating edible or industrially useful oils in commercially significant yields, and include many of the plant species listed above. Such oilseed crops are well known to persons skilled in the art.

RESULTS

Isolation of the TAG1 (DGAT) cDNA from Arabidopsis thaliana

Since one of the most likely defects in AS11 mutant is at the DGAT itself (Table 1; Fig 4), the inventors attempted cloning strategies based on sequence information of enzymes that share common substrates with DGAT. One of the

candidate enzymes that would serve this purpose is the acyl-CoA: cholesterol acyltransferase (ACAT, EC 2.3.1.26) (Chang *et al.*, 1997). Like DGAT, ACAT is an ER protein functioning as an O-acyltransferase by using acyl-CoA as the fatty acyl donor for the esterification of free cholesterol to generate sterol esters. Through a
5 BLAST database search, the inventors identified an *Arabidopsis thaliana* expressed sequence tag (EST) [accession no. AA042298; SEQ ID NO:6] with a deduced amino acid sequence showing 41 % identity to that of the yeast acyl-CoA: cholesterol acyltransferase (Yang *et al.*, 1996, Yu *et al.*, 1996), within the short sequence (104 amino acids) that was available for the EST.

10 The corresponding cDNA (E6B2T7) clone was obtained from the Arabidopsis Biological Resource Center, Columbus, Ohio. Upon complete sequencing, the 878 bp E6B2T7 clone was found to be a partial cDNA. However, the ORF prediction from this partial cDNA confirmed the initial EST search results in that the encoded product is structurally similar to ACAT, especially in the regions at
15 the C-terminus. The inventors were confident that the cDNA contained the 3' untranslated region through an ORF search, although the polyA tail was missing.

The inventors further used the partial cDNA sequence to search against *Arabidopsis thaliana* genomic sequence information. An *Arabidopsis* 'IGF' BAC clone 'F27F23' [accession no. AC003058] was identified to include a region that
20 matched the cDNA, and therefore it was concluded that this was the region encompassing the corresponding gene. Moreover, this BAC clone 'F27F23', is contained in the YAC clone, C1C06E08, which, according to the published map position (http://weeds.mgh.harvard.edu/goodman/c2_b.html), represents a region between centimorgan 35.9 and centimorgan 38.7 on chromosome II; this position is
25 similar to the estimated location for *TAG1*, and the lesion identified by the mutation in AS11 (Katavic *et al.*, 1995). In view of our previous results on the characterization of the AS11 mutant, the map position of this gene strongly suggested that it may encode a DGAT.

To clone a full-length cDNA, a series of oligonucleotide primers were
30 designed, based on the genomic sequences located at different positions 5' upstream of the region covering the partial cDNA. We used these primers in combination with a primer located at the 3' UTR of the partial cDNA (E6B2T7) to

perform PCR reactions with cDNA phagemid prepared from an *Arabidopsis thaliana* (ecotype Columbia) silique-specific cDNA library (Giraudat et al., 1992) as a template. The longest cDNA amplified was 1904 bp, which we subsequently designated as TAG1, and deposited into the Genbank under accession AJ238008

5 [SEQ ID NO:1]. We believe this cDNA represents a full-length clone because its size is in agreement with that of the transcript detected in the northern blot (see Figure 6 a). The longest open reading frame is flanked by a 134-nucleotide 5' untranslated region and a 203-nucleotide 3' untranslated region. There is an in-frame stop codon (TGA at position nt-43) which is followed by an in-frame ATG at

10 position nt-139. It is thus inferred that the ATG at position nt-139 is the initiation codon.

The primary structure of TAG1 predicts a DGAT-related enzyme

The predicted open reading frame of the TAG1 cDNA encodes for a polypeptide of 520 amino acids with a calculated molecular weight of 58993

15 Daltons. With the BLAST search program (Altschul et al. 1990), it was found that the recently reported mouse diacylglycerol acyltransferase [accession no. AF078752] (Cases et al., 1998) is a protein which showed the highest sequence similarity to the deduced amino acid sequence of TAG1 (Figure 5a). TAG1 was also similar to a human acyl CoA: cholesterol acyltransferase-related enzyme

20 [accession no. AF059202]. The human acyl CoA: cholesterol acyltransferase-related enzyme, also known as ARGP1, is most likely to be a DGAT with no significant ACAT activity, although the true nature of the enzyme awaits further confirmation (Oelkers et al., 1998). The similarity between TAG1 and the mammalian DGAT extends over a region of more than 400 amino acids with a

25 sequence identity of about 41%. A putative diacylglycerol/phorbol ester-binding motif, HKW-X-X-RH-X-Y-X-P, a signature sequence observed to be unique to DGAT while absent in the ACATs (Oelkers et al., 1998), is located at amino acids 414-424 ([SEQ ID NO: 7]; Figure 5 a). This diacylglycerol binding motif is also found in the subsequently published *B. napus* DGAT sequences (Nikyforuk et al, 1999;

30 GenBank /EMBL Accession Nos. AF155224, SEQ ID NO:8; AF164434, SEQ ID NO:9). Among other cloned acyltransferases (e.g. GPATs, LPATs, dihydroxyacetone phosphate acyltransferases) it has been reported that there is an invariant proline in a highly hydrophobic block IV that may participate in acyl-CoA

binding (Lewin et al., 1999). In the TAG1 sequence, the hydrophobic block from residues 221-229 containing an invariant proline at residue 224, might constitute such a motif.

TAG1 showed some sequence similarity to other acyl CoA: cholesterol
5 acyltransferases from a number of species (Chang et al., 1997). However, the similarity is largely confined to the C-terminus and is lower (around 30%) than is the similarity of TAG1 to the mammalian DGAT.

The TAG1 protein has multiple hydrophobic domains (Figure 5b) and an analysis by the PC Gene program predicted that the protein has 5 possible
10 transmembrane segments (amino acids 178-195, 233-253, 363-388, 433-476, 486-507). In the mammalian DGAT, a putative tyrosine phosphorylation motif was observed (Cases et al., 1998), but no apparent tyrosine phosphorylation site could be found in TAG1. However, a visual examination revealed a consensus sequence (X-L²⁰⁰-X-K²⁰²-X-X-S²⁰⁵-X-X-X-V²⁰⁹; SEQ ID NO:10) identified as a targeting motif
15 typical of members of the SnRK1 protein kinase family, with serine residue 205 being the residue for phosphorylation. The SnRK1 (SNF1-related protein kinase-1) proteins are a class of Ser/Thr protein kinases that have been increasingly implicated in the global regulation of carbon metabolism in plants (Halford and Hardie, 1998). This consensus SnRK1 targeting motif is also found in the
20 subsequently published *B. napus* DGAT sequences (Nikyforuk et al, 1999; GenBank /EMBL Accession Nos. AF155224; AF164434). Interestingly, similar SnRK1 targeting motifs could also be identified in the *lyso*-phosphatidic acid acyltransferases (LPATs) from coconut (Knutzon et al., 1995) and meadowfoam (Lassner et al., 1995), respectively.

25 *The TAG1 gene is ubiquitously expressed in Arabidopsis*

Northern blot analyses were performed to investigate the expression profile of the TAG1 gene. Total RNA was extracted from different tissues, including roots, leaves, flowers, developing siliques, young seedlings and germinating seeds. The highest steady-state level accumulation of TAG1 transcript was in RNA isolated
30 from germinating seeds and young seedlings (Figure 6a). TAG1 transcripts were also detected in root, leaf and flower tissues, albeit at lower levels. Surprisingly, the TAG1 gene is expressed in developing siliques at a level that is comparable to that

of other vegetative tissues, but lower than that of germinating seeds and young seedlings. This expression profile in general is not inconsistent with the notion that DGAT is present in all plant tissues capable of TAG biosynthesis (Kwanyuan and Wilson, 1986). It has been shown in a number of plant species, including soybean and safflower, that germinating seeds actively synthesize TAGs (Ichihara and Noda, 1981; Kwanyuan and Wilson, 1986; Wilson and Kwanyuan, 1986). The relatively high level of expression in roots is also consistent with the fact that root plastids are capable of synthesizing large amounts of triacylglycerol (Sparace et al., 1992).

Southern blot hybridization (Southern, 1975) was performed with genomic DNA digested with several restriction enzymes including BglII, EcoRI and HindIII. The *TAG1* gene has no internal BglII and HindIII site, while one internal EcoRI site exists. Our Southern analysis suggested that *TAG1* most likely represents a single copy gene in the *Arabidopsis* genome (Figure 6b).

An insertion mutation is found in the TAG1 gene in mutant AS11

Alignment of the genomic sequence (accession no. AC003058; SEQ ID NO:3) with that of the *TAG1* cDNA [SEQ ID NO:1] revealed that the *TAG1* gene contains 16 exons and 15 introns, spanning a region of about 3.4 kb (Figure 7a). DNA containing the *TAG1* allele from AS11 was PCR-amplified and completely sequenced. The AS11 *TAG1* allele has a 147-bp insertion located at the central region of intron 2. The insertion is a duplication of a segment that is composed of 12 bp from the 3' end of intron 1, the entire sequence of exon 2 (81 bp) and 54 bp from the 5' end of intron 2 (Figure 7a).

In order to rule out the possibility of PCR artifacts, two sets of primers were used to perform further PCR amplifications. Primers A and B (see Experimental Procedures, *Primer Strategy*) located in exons 1 and 3, respectively, amplified a DNA fragment that is about 150 bp longer from AS11 (Figure 7b, lane 2) than that from the wild type (Figure 7b, lane 1). The second pair of primers, C and B (Experimental Procedures), with one to be found in both exon 2 and the insertion segment, and the other located in exon 3, generated two amplified fragments from AS11 (Figure 7b, lane 4), while only one from the wild type (Figure 7b, lane 3). Hence these results confirmed that the insertion mutation the inventors identified

through sequencing, reflected the true nature of the mutation in the *TAG1* gene in the AS11 genome.

The AS11 TAG1 transcript has an 81-bp insertion in its open reading frame

Northern blot analyses indicated that there was no difference in the
 5 expression profiles of the *TAG1* gene, between the AS11 mutant and wild type *A. thaliana*. In order to investigate the effect of the mutation at the transcript level, reverse-transcription PCR (RT-PCR) was performed to amplify the *TAG1* transcript from RNA extracted from germinating seedlings of mutant AS11. Sequencing
 10 analysis revealed that there is an 81-bp insertion composed entirely of exon 2 in the transcript from AS11. The exon 2 in the repeat is properly spliced. The alteration of the transcript thus does not disturb the reading frame. However, this additional exon 2 sequence in the AS11 transcript would result in an altered DGAT protein with the 27 amino acid insertion ¹³¹SHAGLFNLCVVVLI AVNSRLIENLMK¹⁵⁷ [SEQ ID NO:11]. The inventors' data shows that this insertion results in a 40-70% reduction
 15 in DGAT activity throughout seed development (Katavic et al., 1995). The 81 bp insert responsible for reduced DGAT activity in AS11 is visible in the comparison of RT-PCR products (Figure 7b: Compare lane 5 (WT) and lane 6 (AS11).) The DNA aberration observed in the AS11 mutant was unexpected, since ethyl methanesulfonate (EMS) generally causes point mutations. Although we cannot
 20 rule out the possibility that this AS11 mutant was the result of a spontaneous mutation event, EMS-induced deletions and insertions have been reported in other systems (Mogami et al., 1986, Okagaki et al., 1991)

The TAG1 gene insertion in Arabidopsis mutant AS11 affects seed triacylglycerol accumulation, but not sterol ester accumulation in seeds.

25 Because TAG1 also showed some sequence homology to acyl CoA: cholesterol acyltransferases (ACATs) from a number of species (Chang et al., 1997), the inventors compared both triacylglycerol and sterol ester accumulation in seeds of the wild-type *A. thaliana* and AS11 mutant. While the triacylglycerol content and TAG/DAG ratios were reduced in AS11 (i.e. increased proportion of
 30 seed oil DAGs,) in contrast, the proportions of sterol esters in WT and AS11 seeds were similar, at 0.8 and 1% of the total lipid extract, respectively (Table 1). If the

TAG1 lesion affected ACAT-like activity, one might expect a reduction in seed sterol esters, but this was not observed. These results indicated that *TAG1* is not involved in sterol-ester homeogenesis, and thus not an acyl CoA: sterol acyltransferase.

TAG1 expression in yeast.

- 5 The *TAG1* cDNA overexpressed in yeast resulted in a 3.5 to 4-fold increase in microsomal DGAT activity compared to plasmid only (*pYES2*) control transformants (Fig. 8), confirming that the *TAG1* gene product functions as a DGAT. When ¹⁴C18:1-CoA was added to the yeast lysates, sterol esters were also labeled *in vitro* (data not shown), but there was no significant difference in the ¹⁴C-labeled
10 sterol esters produced by lysates from the *pYES2* GAL-induced control and the *pYES2:TAG1* Gal-induced transformant. This confirms that the *TAG1* product does not encode an acyl-CoA: sterol acyltransferase (like ACAT).

Complementation of the A. thaliana AS11 Mutant Line by Transformation with the DGAT cDNA.

- 15 The cloned full-length DGAT cDNA was used as a template for PCR amplification with the primers DGATXbaI (CTAGTCTAGAATGGCGATTTTGGGA; SEQ IN NO: 12) and DGATXhoI (GCGCTCGAGTTTCATGACATCGA; SEQ ID NO:13) to provide new restriction sites on each end of the sequence as described in Experimental Procedures. A 1.6kb fragment was excised by a XbaI/KpnI digestion
20 and ligated into the corresponding sites of the pSE129 vector (provided by Dr. P. Covello, PBI/NRC). pSE129A is a vector derived from the plant transformation vector pRD400 (Datla et al. 1992). The vector pSE129A contains the seed-specific napin promoter and the nos terminator cloned into the EcoRI and HindIII sites of the pRD400 plasmid (Figure 9). Hence in the DGAT-pSE129A construct, the
25 *Arabidopsis* DGAT cDNA is under the control of the napin promoter. The construct integrity was confirmed by sequencing.

- The pSE129A containing the napin:DGATcDNA was introduced into *A. tumefaciens*, used to transform *A. thaliana* mutant AS11, and progeny analyzed as described in Experimental Procedures. A number of T₂ transgenic lines were
30 isolated which complemented the fatty acid mutant phenotype found in AS11 (reduced 20:1 and elevated polyunsaturated C₁₈s), restoring the wild-type seed fatty

acid profile (Figure 10). This finding confirms the nature of the lesion in AS11 and directly ties the AS11 lipid phenotype to this mutation.

Over-Expression of the DGAT cDNA in Wild-Type A. thaliana

The cloned full-length DGAT cDNA was used as a template for PCR
5 amplification with the primers DGATXbaI (CTAGTCTAGAATGGCGATTTTGGGA ; SEQ
ID NO:12) and DGATXhoI (GCGCTCGAGTTTCATGACATCGA; SEQ ID NO:13) to
provide new restriction sites on each end of the sequence as described in
Experimental Procedures. A 1.6kb fragment was excised by a XbaI/KpnI digestion
and ligated into the corresponding sites of the pSE129 vector (provided by Dr. P.
10 Covello, PBI/NRC). pSE129A is a vector derived from the plant transformation
vector pRD400 (Datta et al. 1992). The vector pSE129A contains the seed-specific
napin promoter and the nos terminator cloned into the EcoRI and HindIII sites of the
pRD400 plasmid (Figure 9). Hence in the DGAT-pSE129A construct, the
Arabidopsis DGAT cDNA is under the control of the napin promoter. The construct
15 integrity was confirmed by sequencing.

The pSE129A containing the napin:DGATcDNA was introduced into *A.*
tumefaciens, used to transform wild-type *A. thaliana*, and progeny analyzed as
described in Experimental Procedures. A number of T₂ transgenic lines were
isolated which exhibited an increased oil content (Figure 11) an increased average
20 100-seed weight (Figure 12) and a strong linear correlation between the two traits
(Figure 13).

In terms of fatty acyl composition, wild type lines containing over-expressed
DGAT cDNA showed a decrease in the total saturates, and increases in the
monounsaturates and in the 18:1/[18:2 + 18:3] index, as shown in Table 2 below.
25 Such changes are all towards a 'healthier' oil profile, and can be applied directly to
canola, other oilseeds in the *Brassicaceae* and other edible oil crops to produce
similar oil composition improvements.

Table 2

Fatty acid composition of seed oil from *A. thaliana* non-transformed wild-type controls (WT Con) and three T2 transgenic lines (2', 9 and 11) of wild type transformed with the DGAT cDNA under the control of a napin promoter (napin: DGAT).

Line	Total Saturates ^a Wt %	Monounsaturates ^b Wt %	18:1/[18:2 + 18:3] index ^c
WT Control	15.1 ± 0.1	36.7 ± 0.2	29.9 ± 0.6
2' napin:DGAT	13.4	38.6	34.1
9 napin:DGAT	13.1	39.3	35.6
11 napin:DGAT	13.1	38.3	33.0

^a Includes 16:0, 18:0, 20:0, 24:0

^b includes 18:1, 20:1, 22:1, 24:1

^c ([Wt % 18:1] + [Wt % 18:2 + Wt % 18:3]) × 100

EXPERIMENTAL PROCEDURES

Plant Material

- 5 *Arabidopsis thaliana* ecotype Columbia and mutant AS11 were grown under conditions described previously (Katavic et al., 1995). The *A. thaliana* mutant line AS11 was generated and characterized relative to wild type (WT) *A. thaliana* ecotype Columbia, as described by Katavic et al., (1995); (ATCC NO: PTA-1013).

DNA manipulation

- 10 Standard methods and procedures were used for DNA preparation, plasmid propagation and isolation (Sambrook et al., 1989). Sequencing was conducted on an Applied Biosystems Model 373A DNA Sequencing System using the *Taq* DyeDeoxy™ Terminator Cycle Sequencing Kit (Applied Biosystems, Inc.). The nucleotide and the deduced amino acid sequences were compared with sequences
15 available in databanks using the BLAST program (Altschul et al., 1990).

Southern and Northern analysis

Total RNA was extracted from different tissues at various developmental stages, using the method of Lindstrom and Vodkin (1991). RNA samples were denatured with formaldehyde and separated on 1.2% formaldehyde-agarose gels.

- 5 About 5 µg of total RNA was loaded, and the amount of RNA per lane was calibrated by the ethidium bromide-staining intensity of the rRNA bands. Genomic DNA was isolated, digested with restriction enzymes and a Southern blot analysis was performed according to Sambrook et al. (1989). The *TAG1* DNA probe was ³²P labeled by random-priming according to protocols of the manufacture (BRL).

10 *PCR strategy*

Primers used for the amplification of the *TAG1* gene were as follows:

DGAT1 (AGACACGAATCCCATTCACCGA; SEQ ID NO:14), DGAT2

(AGTGGTGACAACGCAGGGATGATG; SEQ ID NO:15), DGAT3 (ATGGTCGCTCCACATTGTGT;

SEQ ID NO:16), DGAT4 (CATACAATCCCATGACATTTATCA; SEQ ID NO:17). DGAT1

- 15 and DGAT2 amplify the 5' half of the *TAG1* gene and DGAT3 and DGAT4 amplify the 3' end of the *TAG1* gene. Genomic DNA from AS11 was used as template for PCR amplification of the mutant *TAG1* allele using the thermal profile: 94° C 3 min; 40 cycles of 94° C 30 seconds, 62° C 45 seconds, 72° C 1 min; and 72° C 15 min.

To further confirm the mutation, primers A (CGACCGTCGGTCCAGCTCATCGG; [SEQ ID

- 20 NO:18]) and B (GCGGCCAATCTCGCAGCGATCTTG; [SEQ ID NO:19]), as well as primers C

(TAAACAGTAGACTCATCATCG; [SEQ ID NO:20]) and B, were used in pairs, respectively,

to amplify the internal fragment containing the mutation. The primers DGAT1 and

DGAT4 were used for PCR amplification of the cDNA with an *A. thaliana* silique cDNA library as template. Primers A and B were also used in RT- PCR

- 25 amplification of the cDNA fragment encompassing the insertion segment.

Construction of TAG1 Multicopy Vector and Transformation and Characterization of DGAT Expression in Yeast

The *TAG1* cDNA was cloned into pBluescript SK as described (Hadjeb and Berkowitz, 1996). The cDNA was cut out from the vector with KpnI/XbaI, and

- 30 subsequently cloned into the respective sites of the yeast expression vector pYes2 (Invitrogen). The construct was confirmed by sequencing. Constructs with *TAG1*

transcription under the control of the GAL1 promoter released a fragment of approximately 1.9 kb. Because the TAG1 fragment has its own initiating ATG codon, the product expressed is not a fusion protein. As a host for yeast expression, an SLC deletion strain (YMN5 [slc1 Δ 2::LEU2 ura3]) (kindly provided by M.M.

- 5 Nagiec and R.C. Dickson, University of Kentucky, Lexington, KY; Nagiec et al., 1993) was used; we reasoned that in this mutant, the endogenous DAG pool may be lower than in WT yeast, and that this would allow us to maximize the activity from over-expressed TAG1 in the presence of exogenously supplied ^{14}C -DAG during in vitro DGAT assays of transformant lysates. Yeast transformation was
- 10 performed according to Elble (1992). YMN5 transformants containing vector only (pYES2) were used as controls. Single colonies were cultured overnight in 20 mL of SD medium (Synthetic Dextrose medium with glucose and without uracil, as described by Ausubel et al., 1995, Vol. 2, p. 13.1.3) on a rotary shaker (270 rpm) at 28°C. Cells were pelleted from the overnight culture and resuspended in 50 mL of
- 15 medium for induction of expression (SD medium containing galactose and without uracil). Cells were reincubated at 28°C, with shaking at 270 rpm, and harvested after 4-6 hr. GAL-induced yeast transformants were harvested by centrifugation at 5000 rpm for 5 min and resuspended in 100 mM Hepes-NaOH, pH 7.4, containing 1 mM EDTA and 1 mM DTT. Cell lysates were prepared using acid-washed glass
- 20 beads as described by Ausubel et al. (1995). Protein in yeast lysates was measured using the Bradford (1976) assay, protein levels in each lysate were normalized and aliquots (250 μg protein) were assayed for DGAT activity as described below.

Lipid Substrates and DGAT Analyses

- ^{14}C -labeled diolein [1- ^{14}C oleic] (Sp. activity 55 mCi/ mmol) was purchased
- 25 from American Radiolabeled Chemicals (St. Louis, MO). The ^{14}C -labelled *sn*-1,2-diolein isomer was purified by TLC on borate-impregnated plates and emulsified in Hepes buffer the presence of 0.2% Tween-20 as described by Taylor et al., (1991). 20:1-CoA, CoASH, ATP, and all other biochemicals were purchased from Sigma.

- DGAT assays were conducted at pH 7.4, with shaking at 100 rpm in a water
- 30 bath at 30°C for 30-60 min. Assay mixtures (0.5 mL final volume) contained lysate protein (250 μg), 90 mM Hepes-NaOH, 0.5 mM ATP, 0.5 mM CoASH, 1 mM MgCl_2 , 200 μM *sn*-1,2 diolein (sp. activity 2 nC/nmol) in 0.02% Tween 20, and 18 μM 20:1-

CoA as the acyl donor. The ^{14}C -labeled TAGs were isolated by TLC and quantified as described by Taylor et al (1991).

Further Lipid and Sterol Ester Analyses in AS11 and WT:

Total lipid extracts (TLEs), and lipid class analyses in WT and the AS11 mutant were performed as described by Taylor *et al.*, (1991; 1992) and by Katavic *et al.*, (1995). Relative seed oil content was also measured by magic angle sample spinning ^1H -NMR, according to the method of Rutar (1989). Analyses were conducted with 200-seed samples of intact wild-type and AS11 seeds using a Bruker AM wide-bore spectrometer (Bruker Analytische Masstechnik GHBH, Silberstreifen D-76287, Rheinstetten4/Karlsruhe, Germany) operating at 360 MHz. To reduce anisotropic line broadening, the seed sample was rotated at 1 kHz in a zirconium rotor oriented 54.7° to the magnetic field. The integration response for resonances attributable to liquid-like oil were summed and the value for AS11 seed was recorded relative to the response for the WT control seed sample, the latter set at a value of 1.00.

Sterol esters were purified from the TLEs by thin layer chromatography (TLC) on Silica H plates developed in hexane: diethyl ether: formic acid (80:20:2, v/v/v). After elution from the silica H with chloroform: methanol (2:1, v/v), the sterol esters were quantified by saponification followed by methylation of the resulting fatty acids with 3N methanolic-HCl. The fatty acid methyl esters (FAMES) were analyzed by GC as described previously (Taylor et al., 1991). The free sterols released by saponification were also analyzed by GC on a 30 m DB-5 column; GC temperature program : initial temp: 180°C , increasing at $10^\circ\text{C}/\text{min}$ to 300°C and held at this temperature for 15 min. The sterol ester content was reported as a % of the TLE; i.e. FAMES released from sterol esters calculated as proportion of the FAMES released by transmethylation of the total lipid extract (TLE).

Construction of Plant Transformation Vector Containing the Wild-Type DGAT gene for Over-Expression in WT A. thaliana and Complementation of the A. thaliana AS11 Mutant:

Two primers:

Gen 1 (GAGAGGATCCACGCTCACGACCCATTCTTCCCG; [SEQ ID NO:21]), and

Gen 2 (AAGAAGGATCCATCCCCAAAACGGGACCACCAA; [SEQ ID NO:22])

- 5 were synthesized according to sequences upstream and downstream of the TAG1 gene. These primers were used to PCR amplify a genomic fragment of 5.1 kb from wild-type *A. thaliana*. The PCR fragment was purified and digested with BamHI and inserted into the corresponding site in plasmid pRD400 (Datla et al. 1992) to generate the plant transformation vector DGATg-pRD400.

10 *Construction of DGAT cDNA Plant Transformation Vector for Seed-Specific Expression:*

- The cloned full-length DGAT cDNA was used as a template for PCR amplification with the primers DGATXbaI (CTAGTCTAGAATGGCGATTTTGA; SEQ ID NO:12) and DGATXhoI (GCGCTCGAGTTTCATGACATCGA; SEQ ID NO: 13) to
 15 provide new restriction sites on each end of the sequence. The PCR profile was as follows: 94°C 1 min; 30 cycles of 94°C 30 seconds, 55°C 30 seconds, 72°C 1 min; and 72°C 5 min. The PCR product was then ligated into the PCR-2.1 vector (InVitrogen). A 1.6kb fragment was excised by a XbaI/KpnI digestion and ligated into the corresponding sites of the pSE129 vector (provided by Dr. P. Covello,
 20 PBI/NRC). pSE129A is a vector derived from the plant transformation vector pRD400 (Datla et al. 1992). The vector pSE129A contains the seed-specific napin promoter and the nos terminator cloned into the EcoRI and HindIII sites of the pRD400 plasmid (See Figure 9). Hence in the DGAT-pSE129A construct, the *Arabidopsis* DGAT cDNA is under the control of the napin promoter. The construct
 25 integrity was confirmed by sequencing.

Transformation of Agrobacterium with Plant DGAT Vector Constructs:

- Electrocompetent *Agrobacterium* cells, GV3101 (pMP90) strain, were prepared as follows: An *Agrobacterium* culture was grown 24 to 48 hrs in 2YT, and when the absorbance at 600 nm reached 0.5 to 0.7, the cells were chilled on ice
 30 and pelleted by centrifugation (5,000 x g, 10 min in a GSA rotor at 4°C). The pellet was washed in 1, 0.5 and 0.02 volumes of cold 10% sterile glycerol and

resuspended in 0.01 volume of cold 10% glycerol. The electrocompetent cells were then frozen in liquid N₂ and stored at -70°C. The *Agrobacterium* cells were transformed by electroporation with 20-50 ng of transforming DNA (either DGATg-pRD400 or DGAT-pSE129A) according to the manufacturer's instructions, plated on
5 a selective medium (LB with 50 µg/mL kanamycin) and incubated overnight at 28°C. Single transformed cells were grown overnight (28°C, 225 r.p.m.) in 5 mL LB with 50 µg/mL Kanamycin and 25 µg/mL Gentamycin. DNA extraction and purification were performed with a Qiaprep Spin Miniprep kit (Qiagen). The fidelity of the construct was re-checked by DNA sequencing before plant transformation.

10 *Transformation of Arabidopsis thaliana:*

The transformation protocol was adapted from that described by Clough and Bent (1998). Seeds of *Arabidopsis thaliana* ecotype Columbia and mutant AS11 (Katavic *et al.*, 1995) were grown at 22°C under fluorescent illumination (120 µE·m⁻²·s⁻¹) in a 16 h light/8 hour dark regime. Typically, four to six plants were raised in a
15 10 cm² pot in moistened Terra-lite Redi-earth (W. R. Grace & Co. Canada Ltd. Ajax, ON, Canada). To prevent the soil mix in the pot from falling into the inoculation media, soil was mounded as a platform with seeds sown on top, and the whole pot covered by a nylon window screen and secured by a rubber band. Plants were vacuum infiltrated in an *Agrobacterium* suspension when the first flowers started
20 opening.

To grow *Agrobacterium*, a 5 mL suspension in LB medium containing 50 µg/mL kanamycin and 25 µg/mL gentamycin was cultured overnight at 28°C. The day before infiltration, this "seed culture" was divided into four flasks containing 250 mL of LB medium supplemented with 50 µg/mL kanamycin and 25 µg/mL
25 gentamycin. These culture were grown overnight at 28°C. The next morning after the absorbance at 600 nm was checked (approximately = 1.0), the cells were harvested by centrifugation (5,000 x g, 10 min in a GSA rotor at room temperature) and resuspended in the infiltration medium (sucrose 5%; Silwet-77 0.005% in water) to obtain an optical density at 600 nm of 0.8. The *Agrobacterium* suspension was
30 then poured into a beaker and the potted plants inverted into the beaker so that the flowers and bolts were submerged. The beaker was then placed into a large Bell jar and a vacuum drawn using a vacuum pump, until bubbles formed on the stem

surfaces and the solution started to bubble slightly, and then the vacuum was released rapidly. [Note: The necessary time and pressure will vary from one lab setup to the next, but good infiltration is visibly apparent as uniformly darkened, water-soaked tissue.] Pots were removed from the beaker, laid on their side in a plastic tray and covered with a plastic dome, to maintain humidity. The following day, the plants were uncovered, set upright and allowed to grow for approximately four weeks in a growth chamber under continuous light conditions as described by Katavic et al., (1995). When the siliques were mature and dry, seeds were harvested and selected for positive transformants.

10 *Selection of Putative Transformants (Transgenic plants) and Analysis of Transgenic Plants:*

For each construct, seeds were harvested in bulk. Seeds were surface-sterilized by submerging them in a solution containing 20% bleach and 0.01% Triton X-100 for 20 min, followed by three rinses with sterile water. Sterilized seeds were then plated by resuspending them in sterile 0.1% phytagar at room temperature (about 1 mL phytagar for every 500-1000 seeds), and then applying a volume containing 2,000-4,000 seeds onto 150 x 15 mm kanamycin selection plate. Plates were incubated for 2 days in the cold without light, and then grown for 7-10 days in a controlled environment (22°C under fluorescent illumination ($120 \mu\text{E}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) in a 16 h light/8 hour dark regime). The selection media contain $\frac{1}{2}$ MSG medium, 0.8% phytagar, 3% sucrose, 50ug/mL kanamycin and 50ug/ mL Timentin. Petri dishes and lids were sealed with a Micropore™ surgical tape tape (3M Canada, London, ON, Canada). After 7-10 days, drug-resistant plants that had green leaves and well established roots within the medium were identified as transformants and at the 3-5 leaf stage, selected transformants were transplanted into flats filled with heavily moistened soil mix. Transformants were grown to maturity and mature seeds (T_2 generation as defined in Katavic et al., (1994)) were harvested from individual plants, and further analyzed.

DNA isolation from and analysis of Transformants

30 Genomic DNA was isolated from individual T_1 plants following the protocol of Dellaporta et al., (1983). A PCR amplification using the paired primers described previously for the DGAT cDNA or for the DGAT gene, was performed to confirm the

presence of the cDNA or the gene, respectively, in the T₁ transformants. Southern analyses (Southern, 1975) were performed to select the transformants containing a single copy of the inserted fragment. DNA samples were digested with restriction enzymes (Bgl II for the DGAT cDNA and Eco RI for the DGAT gene), resolved by

5 electrophoresis on a 1% agarose gel, and Southern blotting performed using a nylon filter (Hybond-N+, Amersham) according to Sambrook *et al.* (1989). The DGAT cDNA fragment, labelled with α -[³²P] dCTP (NEN/DuPont) using the Random Primer DNA labelling kit (Gibco BRL), was used as a probe. Hybridization was performed at 60°C according to Church and Gilbert (1984). The filter was then

10 exposed to Kodak X-OMAT-AR film.

DEPOSIT INFORMATION

The following biological material has been deposited at the American Type Culture Collection (ATCC) of 10801 University Boulevard, Manassas, Virginia, 20110-2209, U.S.A. All of these deposits were made on behalf of the

15 Applicant/Assignee (National Research Council of Canada) under the terms of the Budapest Treaty on the dates indicated, and have been given the accession numbers shown below.

	Deposited Material	Date of Deposit	Accession No.
	Arabidopsis DGAT gene	November 29, 1999	PTA-988
20	Arabidopsis DGAT cDNA	November 29, 1999	PTA-989
	Arabidopsis ASII seeds	December 3, 1999	PTA-1013

The deposit receipts are shown later in this description.

25 SEQUENCE LISTING FREE TEXT

The Sequence Listing provided below contains free text entries in respect of SEQ ID NOs: 12 to 22. The free text used in the Sequence Listing is repeated as follows:

	SEQ ID NO:12	Primer of DGATXbal
30	SEQ ID NO:13	Primer of DGATXhol

	SEQ ID NO:14	Primer DGAT1
	SEQ ID NO:15	Primer DGAT2
	SEQ ID NO:16	Primer DGAT3
	SEQ ID NO:17	Primer DGAT 4
5	SEQ ID NO:18	Primer A
	SEQ ID NO:19	Primer B
	SEQ ID NO:20	Primer C
	SEQ ID NO:21	Primer Gen 1
	SEQ ID NO:22	Primer Gen 2.

10

A summary of all of the listed sequences is provided below for ease of review:

SEQ ID NO:1 - (pDGAT; vector containing isolated and purified deoxyribonucleic acid cDNA; ATCC No PTA-989), Genbank/EMBL Accession No. AJ238008.

- 15 SEQ ID NO:2 - The deduced amino acid sequence of the *Arabidopsis* DGAT (AtTAG1) protein.

SEQ ID NO:3 - (pgenomic DGAT; vector containing isolated and purified genomic deoxyribonucleic acid (genomic DNA) ATCC No PTA-988).

- 20 SEQ ID NO:4 - MDGAT, mouse DGAT [GenBank/EMBL Accession No. AF078752 (Cases *et al.*, 1998)].

SEQ ID NO:5 - HARGP1, human ARGPI protein [GenBank/EMBL Accession No. AF059202; Oelkers *et al.*, 1998].

SEQ NO:6 - *Arabidopsis thaliana* expressed sequence tag (EST) [accession no. AA042298).

- 25 SEQ ID NO:7 - A diacylglycerol/phorbol ester-binding motif found in SEQ ID NO:2, SEQ ID NO:8 and SEQ ID NO:9 (⁴¹⁴HKWMVRHIYFP⁴²⁴).

SEQ ID NO:8 - *B. napus* DGAT amino acid sequence GenBank EMBL Accession No AF155224.

SEQ ID NO:9 - *B. napus* DGAT amino acid sequence GenBank/EMBL Accession
No. AF164434.

5 SEQ ID NO: 10 - Targeting motif typical of members of the SnRK1 protein kinase
family found in SEQ ID NO:2, SEQ ID NO:8 and SEQ ID NO:9 X-L²⁰⁰-X-K²⁰²-
X-X-S²⁰⁵-X-X-X-V²⁰⁹

SEQ ID NO:11 - A 27 amino acid insertion repeat in SEQ ID NO:2 found in the
Arabidopsis thaliana AS11 mutant.

¹³¹SHAGLFNLCVVVLIAVNSRLIENLMK¹⁵⁷

10 SEQ IN NO: 12 - CTAGTCTAGAATGGCGATTTTGA (nucleotide sequence of
Primer DGATXbal).

SEQ ID NO:13 - GCGCTCGAGTTTCATGACATCGA (nucleotide sequence of
Primer DGATXhol).

SEQ ID NO:14 - AGACACGAATCCCATTCACCGA (nucleotide sequence of
Primer DGAT1).

15 SEQ ID NO:15 - AGTGGTGACAACGCAGGGATGATG (nucleotide sequence of
Primer DGAT2).

SEQ ID NO:16 - ATGGTCGCTCCACATTGTGT (nucleotide sequence of Primer
DGAT3).

20 SEQ ID NO:17 - CATACAATCCCCATGACATTTATCA (nucleotide sequence of
Primer DGAT4).

SEQ ID NO:18 - CGACCGTCGGTTCAGCTCATCGG (nucleotide sequence of
Primer A).

SEQ ID NO:19 - GCGGCCAATCTCGCAGCGATCTTG (nucleotide sequence of
Primer B).

25 SEQ ID NO:20 - TAAACAGTAGACTCATCATCG (nucleotide sequence of
Primer C).

SEQ ID NO:21 - GAGAGGATCCACGCTCACGACCCATTCTTCCCG (nucleotide sequence of primer Gen 1).

SEQ ID NO:22 - AAGAAGGATCCATCCCCAAAACGGGACCACCAA

(nucleotide sequence of primer Gen 2).

5 SEQ ID NO:23 - AS11 mutant DGAT cDNA nucleotide sequence.

SEQ ID NO:24 - AS11 mutant DGAT genomic DNA nucleotide sequence.

SEQ ID NO:25 - the deduced amino acid sequence of SEQ ID NO:23.

ATCC

10801 University Blvd • Manassas, VA 20110-2209 • Telephone: 703-365-2700 • FAX: 703-

BUDAPEST TREATY ON THE INTERNATIONAL RECOGNITION OF
THE DEPOSIT OF MICROORGANISMS FOR THE PURPOSES OF PATENT PROCEDURE**INTERNATIONAL FORM**RECEIPT IN THE CASE OF AN ORIGINAL DEPOSIT ISSUED PURSUANT TO RULE 7.3
AND VIABILITY STATEMENT ISSUED PURSUANT TO RULE 10.2

To: (Name and Address of Depositor or Attorney)

NRC Canada
Attn: David C. Taylor
Plant Biotechnology Institute
110 Gymnasium Place
Saskatoon, SK S7N 0W9 Canada**RECEIVED**
KEGB**DEC 10 1999**

ENTERED _____

CHECKED _____

Deposited on Behalf of: National Research Council of Canada

Identification Reference by Depositor:

Patent Deposit Designation

Arabidopsis DGAT gene
Arabidopsis DGAT cDNAPTA-988
PTA-989The deposits were accompanied by: ___ a scientific description X a proposed taxonomic description indicated above. The deposits were received November 29, 1999 by this International Depository Authority and have been accepted.AT YOUR REQUEST: X We will not inform you of requests for the strains.

The strains will be made available if a patent office signatory to the Budapest Treaty certifies one's right to receive, or if a U.S. Patent is issued citing the strains, and ATCC is instructed by the United States Patent & Trademark Office or the depositor to release said strains.

If the cultures should die or be destroyed during the effective term of the deposit, it shall be your responsibility to replace them with living cultures of the same.

The strains will be maintained for a period of at least 30 years from date of deposit, or five years after the most recent request for a sample, whichever is longer. The United States and many other countries are signatory to the Budapest Treaty.

The viability of the cultures cited above was tested December 2, 1999 On that date, the cultures were viable.

International Depository Authority: American Type Culture Collection, Manassas, VA 20110-2209 USA.

Signature of person having authority to represent ATCC:


Barbara M. Halley, Administrator, Patent Depository

Date: December 3, 1999

cc: Kirby, Eades, Gale and Baker

ATCC

10801 University Blvd • Manassas, VA 20110-2209 • Telephone: 703-365-2700 • FAX: 703-

**BUDAPEST TREATY ON THE INTERNATIONAL RECOGNITION OF
THE DEPOSIT OF MICROORGANISMS FOR THE PURPOSES OF PATENT PROCEDURE****INTERNATIONAL FORM****RECEIPT IN THE CASE OF AN ORIGINAL DEPOSIT ISSUED PURSUANT TO RULE 7.3
AND VIABILITY STATEMENT ISSUED PURSUANT TO RULE 10.2****To: (Name and Address of Depositor or Attorney)**

National Research Council of Canada
 Attn: David C. Taylor
 Plant Biotechnology Institute
 110 Gymnasium Place
 Saskatoon, SK S7N 0W9
 Canada

Deposited on Behalf of: National Research Council of Canada**Identification Reference by Depositor:****Patent Deposit Designation***Arabidopsis* AS11

PTA-1013

The seeds were accompanied by: a scientific description ☒ a proposed taxonomic description indicated above.
 The seeds were received December 3, 1999 by this International Depository Authority and have been accepted.

AT YOUR REQUEST: ☒ We will not inform you of requests for the seeds..

The seeds will be made available if a patent office signatory to the Budapest Treaty certifies one's right to receive,
 or if a U.S. Patent is issued citing the seeds and ATCC is instructed by the United States Patent & Trademark Office
 or the depositor to release said seeds.

If the seeds should die or be destroyed during the effective term of the deposit, it shall be your responsibility to
 replace them with viable seeds of the same.

The seeds will be maintained for a period of at least 30 years from date of deposit, or five years after the most recent
 request for a sample, whichever is longer. The United States and many other countries are signatory to the Budapest
 Treaty.

The viability of the seeds cited above was tested December 23, 1999. On that date, the seeds were viable.

International Depository Authority: American Type Culture Collection, Manassas, VA 20110-2209 USA.

Signature of person having authority to represent ATCC:

Barbara M. Hailey
 Barbara M. Hailey, Administrator, Patent Depository

Date: January 3, 2000

cc: Kirby, Eades et al.

SUBSTITUTE SHEET (RULE 28)

096354-032901

REFERENCES RELEVANT TO THE CURRENT INVENTION

- Altschul, S.F., Gish, W., Miller, W., Myers, E.W. and Lipman, D.J (1990) Basic local alignment search tool. *J. Mol. Biol.* 215, 403-410.
- Ausubel, F.M., Brent, R., Kingston, R.E., Moore, D.D., Seidman, J.G., Smith, J.A., and Stuhl, K., eds (1995). *Current Protocols in Molecular Biology*, Vols 1, 2, and 3. Wiley, New York.
- Bechtold, N., Ellis, J., and Pelletier, G. (1993) *In planta Agrobacterium-mediated gene transfer by infiltration of adult Arabidopsis thaliana plants. C R Acad Sci Paris, Sciences de la vie/Life sciences* 316: 1194-1199.
- Becker, D., Brettschneider, R. and Lörz, H. (1994) Fertile transgenic wheat from microprojectile bombardment of scutellar tissue. *Plant J.* 5: 299-307.
- Bernerth R, Frentzen M (1990) Utilization of erucoyl-CoA by acyltransferases from developing seeds of *Brassica napus* (L.) involved in triacylglycerol biosynthesis. *Plant Sci* 67: 21-28.
- Bradford M.M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72, 248-254.
- Budziszewski, G.J., Croft, K.P.C. and Hildebrand, D.F. (1996) Uses of biotechnology in modifying plant lipids. *Lipids* 31: 557-569.
- Cao, Y-Z and Huang, AHC (1986) Diacylglycerol acyltransferase in maturing oil seeds of maize and other species. *Plant Physiol.* 82: 813-820.
- Cao Y-Z, Huang AHC (1987) Acyle coenzyme A preference of diacylglycerol acyltransferase from maturing seeds of *Cuphea*, maize, rapeseed and canola. *Plant Physiol.* 84: 762-765
- Cases, S., Smith, J.S., Zheng, Y-W., Myers, H.M., Lear, S.R., Sande, E., Novak, S., Collins, C., Welch, C.B., Lusi, A.J., Erickson, S.K. and Farese, R.V., JR. (1998) Identification of a gene encoding a acyl CoA: diacylglycerol acyltransferase, a key enzyme in triacylglycerol synthesis. *Proc. Nat'l. Acad. Sci. USA*, 95, 13018-13023.

- Chang, T.Y., Chang, C.C.Y. and Cheng, D. (1997) Acyl-Coenzyme A: Cholesterol Acyltransferase. *Annu. Rev. Biochem.* 66, 613-38.
- Church, G.M. and Gilbert, W. (1984) Genomic sequencing. *Proc. Natl. Acad. Sci. USA.* 81, 1991-95.
- Clough, S.J. and Bent, A.F. (1998) Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *The Plant Journal* 16, 735-43
- Datla, R.S.S., Hammerlindl, J.K., Panchuk, B., Pelcher, L.E. and Keller, W.A. (1992) Modified binary plant transformation vectors with the wild-type gene encoding NPTII. *Gene* 211: 383-384.
- Datla, R.S.S., Bekkaoui, F., Hammerlindl, J., Pilate, G., Dunstan, D.I. and Crosby, W.L. (1993) Improved high-level constitutive foreign gene expression in plants using an AMV RNA4 untranslated leader sequence. *Plant Sci.* 94: 139-149.
- Datla, R., Anderson, J.W. and Selvaraj, G. (1997) Plant promoters for transgene expression. *Biotechnology Annual Review* 3: 269-296.
- DeBlock, M., DeBrouwer, D. and Tenning P. (1989) Transformation of *Brassica napus* and *Brassica oleracea* using *Agrobacterium tumefaciens* and the expression of the *bar* and *neo* genes in the transgenic plants. *Plant Physiol.* 91: 694-701.
- De Lange, P., Van Blokland, R., Kooter, J.M. and Mol, J.M.N. (1995) Suppression of flavenoid flower pigmentation genes in *Petunia hybrida* by the introduction of antisense and sense genes. *In: Gene Silencing in Higher Plants and Related Phenomena in Other Eukaryotes*. P.Meyer (Ed.), Springer-Verlag, Berlin, pp. 55-75.
- Dellaporta, S.L., Wood, J. and Hicks, J.B. (1983) A plant DNA miniprep: Version II. *Plant Mol. Biol. Rep.* 1, 19-21
- Elble, R. (1992). A simple and efficient procedure for transformation of yeasts. *Biotechniques* 13,18-20.
- Focks, N. and Benning, C. (1998) *wrinkled1*: A novel, low-seed-oil mutant of *Arabidopsis* with a deficiency in the seed-specific regulation of carbohydrate metabolism. *Plant Physiol.* 118, 91-101.

- Frentzen M (1993) Acyltransferases and triacylglycerols. *In*: Moore, Jr. TS, editor, *Lipid Metabolism in Plants*, pp. 195-230. CRC Press, Ann Arbor,.
- Giraudat, J., Hauge, B.M., Vaion, C., Smalle, J., Parcy, F., Goodman, H.M. (1992) Isolation of the *Arabidopsis* ABI3 gene by positional cloning. *Plant Cell* 4, 1251-1261.
- Goodman, H.M., Ecker, J.R. and Dean, C. (1995) The genome of *Arabidopsis thaliana*. *Proc. Nat'l. Acad. Sci. USA* 92: 10831-10835.
- Hadjeb N and Berkowitz GA (1996) Preparation of T-over-hang vectors with high PCR product cloning efficiency. *Biotechniques* 20: 21-22.
- Halford, N.G. and Hardie, D.G. (1998) SNF1-related protein kinases: global regulators of carbon metabolism in plants? *Plant Mol. Biol.* 37, 735-748.
- Hitz, W.D., Mauvis, C.J., Ripp, K.G., Reiter, R.J., DeBonte, L. and Chen, Z. (1995) The use of cloned rapeseed genes for cytoplasmic fatty acid desaturases and the plastid acyl-ACP thioesterases to alter relative levels of polyunsaturated and saturated fatty acids in rapeseed oil. *Proc. 9th Internat'nal Cambridge Rapeseed Congress UK*, pp. 470-472.
- Ichihara, K. and Noda, M. (1981) Lipid synthesis in germinating safflower seeds and protoplasts. *Phytochemistry* 20, 1245-1249.
- Ichihara, K., Takahashi, T. and Fujii, S. (1988) Diacylglycerol acyltransferase in maturing safflower seeds: its influences on the fatty acid composition of the triacylglycerol and on the rate of triacylglycerol synthesis. *Biochim. Biophys. Acta* 958, 125-129.
- Jorgensen, R.A. and Napoli, C.A. (1994) Genetic engineering of novel plant phenotypes. U.S. Patent No. 5283184.
- Josefsson, L-G, Lenman M, Ericson ML and Rask L (1987) Structure of a gene encoding the 1.7S storage protein, napin, from *Brassica napus*. *J Biol Chem* 262: 12196-12201.

- Katavic, V., Haughn, G.W., Reed, D., Martin, M. and Kunst, L. (1994) In planta transformation of *Arabidopsis thaliana*. *Mol. Gen. Genet.* 245: 363-370.
- Katavic, V., Reed, D.W., Taylor, D.C., Giblin, E.M., Barion, D.L., Zou, J-T., MacKenzie, S.L., Covello, P.S. and Kunst, L. (1995) Alteration of fatty acid composition by an EMS-induced mutation in *Arabidopsis thaliana* affecting diacylglycerol acyltransferase activity. *Plant Physiol.* 108, 399-409.
- Kinney, A.J. (1995) Improving soybean seed quality. In: *Induced Mutations and Molecular Techniques for Crop Improvement*. International Atomic Energy Agency, Vienna, Austria., pp. 101-113.
- Kinney, A.J. (1997) Genetic engineering of oilseeds for desired traits. In: *Genetic Engineering*, Vol. 19 (J.K. Setlow, ed.), Plenum Press, NY., pp. 149-166.
- Kishore G.M. and Somerville, C.R. (1993) Genetic engineering of commercially useful biosynthetic pathways in transgenic plants. *Current Opinion in Biotechnology.* 4: 152-158.
- Knutzon, D.S., Thompson, G.A., Radke, S.E., Johnson, W.B., Knauf, V.C., and Kridl, J.C. (1992) Modification of Brassica seed oil by anti-sense expression of a stearyl-acyl carrier protein desaturase gene. *Proc. Nat'l Acad. Sci. USA*, 89: 2624-2628.
- Knutzon, D.S., Lardizabal, K.D., Nelson, J.S., Bleibaum, J.L., Davis, H.M and Metz, J. (1995) Cloning of a coconut endosperm cDNA encoding a 1-acyl-sn-glycerol-3-phosphate acyltransferase that accepts medium chain length substrates. *Plant Physiol.* 109, 999-1006.
- Kwanyuen, P. and Wilson, R.F. (1986) Isolation and purification of diacylglycerol acyltransferase from germinating soybean cotyledons. *Biochim. Biophys. Acta* 877, 238-245.
- Lacey DJ, Hills MJ (1996) Heterogeneity of the endoplasmic reticulum with respect to lipid synthesis in developing seeds of *Brassica napus* L. *Planta* 199: 545-551.
- Lagercrantz, U., Putterill, J., Coupland, G. and Lydiate, D. (1996) Comparative mapping in *Arabidopsis* and *Brassica*, fine scale genome collinearity and congruence of genes controlling flowering. *Plant J.* 9: 13-20.

- Lassner, M.W., Levering, C.K., Davis, H.M. and Knutzon, D.S. (1995)
Lysophosphatidic acid acyltransferase from meadowfoam mediates insertion of
erucic acid at the sn-2 position of triacylglycerol in transgenic rapeseed oil. *Plant
Physiol.* 109, 1389-1394.
- Lassner, M.W., Lardizabal, K, and Metz, J.G. (1996) A jojoba β -ketoacyl-CoA synthase
cDNA complements the canola fatty acid elongation mutation in transgenic plants.
The Plant Cell, 8: 281-292.
- Lewin TM, Wang P and Coleman RA (1999) Analysis of amino acid motifs diagnostic
for the sn-glycerol-3-phosphate acyltransferase reaction. *Biochemistry* 38: 5764-
5771.
- Lindstrom J.T. and Vodkin L.O. (1991) A soybean cell wall protein is affected by seed
colour genotype. *Plant Cell* 3:561-571
- Little D, Weselake RJ, Pomeroy MK, Furukawa-Stoffer T and Bagu J (1994)
Solubilization and characterization of diacylglycerol acyltransferase from microspore-
derived cultures of oilseed rape. *Biochem J* 304: 951-958.
- Lloyd, A.M., Walbot, V. and Davis, R.W. (1992) Arabidopsis and Nicotiana anthocyanin
production activated by maize regulators R and C1. *Science* 258: 1773-1775.
- MacKenzie, S.L. and Jain, R.K. (1997) Improvement of oils crops via biotechnology.
Recent Res. Dev. In Oil Chem. 1: 149-158.
- Mayorek, N, Grinstein I, and Bar-Tana J (1989) Triacylglycerol synthesis in cultured rat
hepatocytes. The rate-limiting role of diacylglycerol acyltransferase. *Eur J Biochem*
182: 395-400.
- Meyer, P. (1995) Understanding and controlling transgene expression. *Trends in
Biotechnology*, 13: 332-337.
- Meyerowitz, E.M. (1987) Arabidopsis thaliana. *Ann. Rev. Genet.* 21: 93-111.
- Meyerowitz, E.M. and Chang, C. (1985) Molecular biology of plant growth and
development: Arabidopsis thaliana as an experimental system. In: *Developmental
Biology*, Vol. 5, Plenum Press, NY., pp. 353-366.

- Mogami, K., O'Donnell, P.T., Bernstein, S.I., Wright, T.R.F and Emerson, C.P., JR. (1986) Mutations of the *Drosophila* myosin heavy-chain gene: effects on transcription, myosin accumulation, and muscle function. *Proc. Nat'l. Acad. Sci. USA.* 83, 1393-1397.
- Mol, J.M.N., Van der Krol, A.R., Van Tunen, A.J., Van Blokland, R., De Lange, P. and Stuitje, A.R. (1990) Regulation of plant gene expression by antisense RNA. *FEBS Lett.* 268: 427-430.
- Moloney, M.M., Walker, J.M. and Sharma, K.K. (1989) High efficiency transformation of *Brassica napus* using *Agrobacterium* vectors. *Plant Cell Rep. B:* 238-242.
- Nagiec, M.M., Wells, G.B., Lester, R.L., and Dickson, R.C. (1993). A suppressor gene that enables *Saccharomyces cerevisiae* to grow without making sphingolipids encodes a protein that resembles an *Escherichia coli* fatty acyltransferase. *J. Biol. Chem.* 268, 22156-22163.
- Nehra, N.S., Chibbar, R.N., Leung, N., Caswell, K., Mallard, C., Steinhauer, L. Baga, M. and Kartha K.K. (1994) Self-fertile transgenic wheat plants regenerated from isolated scutellar tissues following microprojectile bombardment with two distinct gene constructs. *Plant J* 5: 295-297.
- Nykiforuk C, Laroche A and Weselake RJ (1999) Isolation and sequence analysis of a novel cDNA encoding a putative diacylglycerol acyltransferase from a microspore-derived cell suspension culture of *Brassica napus* L. cv Jet Neuf (Accession No. AF155224.), *Plant Physiology* 120: 1207.
- Oelkers, P., Behar, A., Cromley, D., Bilthelmer, J.T. and Sturley, S.T. (1998) Characterization of two human genes encoding acyl Coenzyme A: cholesterol acyltransferase-related enzymes. *J. Biol. Chem.* 273, 26765-26771.
- Okagaki, R.J., Neuffer, M.G. and Wessler, S.R. (1991) A deletion common to two independently derived Waxy mutations in maize. *Genetics* 128, 425-431.
- Okuley, J., Lightner, J., Feldmann, K., Yadav, N., Lark, E. and Browee, J. (1994) *Arabidopsis fad2* gene encodes the enzyme that is essential for polyunsaturated lipid synthesis. *The Plant Cell* 6: 147-158.

- Perry, H.Y. and Harwood, J.L. (1993a) Changes in the lipid content of developing seeds of *Brassica napus*. *Phytochemistry* 32: 1411-1415.
- Perry, H.Y. and Harwood, J.L. (1993b) Use of [2-3H] glycerol precursor in radiolabelling studies of acyl lipids in developing seeds of *Brassica napus*. *Phytochemistry* 34: 69-73.
- Poirier, Y., Dennis DE, Klomparens K and Somerville C (1992) Polyhydroxybutyrate, a biodegradable thermoplastic produced in transgenic plants. *Science* 256: 520-523.
- Poirier, Y., Nawrath C and Somerville C (1995) Production of polyhydroxyalkanoates, a family of biodegradeable plastics and elastomers in bacteria and plants butyrate, a biodegradable thermoplastic produced in transgenic plants. *Bio-Technology*, 13: 142-150.
- Poirier, Y., Ventre, G and Caldelari, D(1999) Increased flow of fatty acids toward β oxidation in developing seeds of *Arabidopsis* deficient in diacylglycerol acyltransferase activity or synthesizing medium-chain-length fatty acids. *Plant Physiology* 121: 1359-1366.
- Potrykus, I. (1991) Gene transfer to plants: Assessment of published approaches and results. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 42: 205-225.
- Radke SE, Andrews BM, Moloney MM, Crouch ML, Kridl JC, and Knauf VC (1988) Transformation of *Brassica napus* L. using *Agrobacterium tumefaciens*: developmentally regulated expression of a reintroduced napin gene. *Theor. Appl. Genet.* 75: 685-694.
- Rhodes, C.A., Pierce, D.A., Mettler, I.J., Mascarenhas, D. and Detmer, J.J. (1988) Genetically transformed maize plants from protoplasts. *Science* 240: 204-207.
- Rutar, V. (1989) Magic angle sample spinning NMR spectroscopy of liquids as a non-destructive method for studies of plant seeds. *J. Agric. Food. Chem.* 37, 67-70.
- Sambrook J., Fritsch E.F. and Maniatis T. (1989) In *Molecular Cloning, A Laboratory Manual*, 2nd edition. Cold Spring Harbor Laboratory Press.

- Sanford, J.C., Klein, T.M., Wolf, E.D. and Allen, N. (1987) Delivery of substances into cells and tissues using a particle bombardment process. *J. Part. Sci. Technol.* 5: 27-37.
- Settlage, SH, Wilson RF and Kwanyuen, P. (1995) Localization of diacylglycerol acyltransferase to oil body associated endoplasmic reticulum. *Plant Physiol. Biochem.* 33: 399-407.
- Shimamoto, K., Terada, R., Izawa, T. and Fujimoto, H. (1989) Fertile transgenic rice plants regenerated from transformed protoplasts. *Nature* 338: 274-276.
- Somerville, C.R. (1993) Future prospects for genetic modification of the composition of edible oils from higher plants. *Am. J. Clin. Nutr.* 58 (2 Suppl.): 270S-275S.
- Songstad, D.D., Somers, D.A. and Griesbach, R.J. (1995) Advances in alternative DNA delivery techniques. *Plant Cell, Tissue and Organ Culture* 40: 1-15.
- Southern E.M. (1975) Detection of specific sequences among DNA fragments separated by gele electrophoresis. *J. Mol. Biol.* 98: 503-517.
- Sparace S.A., Kleppinger-Sparace K.F., Stahl R.J., Xue L. and Qi, Q. (1992) Lipid biosynthesis in pea root plastids and some effects of glycolytic intermediates. In SL MacKenzie and DC Taylor, eds. *Seed Oils for the Future*, AOCS Press, Champaign, IL. pp 52-60.
- Stark, D.M., Timmerman, K.P., Barry, G.F., Preiss, J. and Kishore, G.M. (1992) Regulation of the amount of starch in plant tissues by ADP glucose pyrophosphorylase. *Science* 258: 287-292.
- Stobart AK, Stymne S, Höglund S. (1986) Safflower microsomes catalyse oil accumulation in vitro: A model system. *Planta* 169: 33-37.
- Stymne, S. and Stobart, A.K. (1987) Triacylglycerol Biosynthesis. In Stumpf, P.K. ed, *The Biochemistry of Plants*, Academic Press, New York. 9, 175-214.
- Taylor, CB. (1998) Comprehending cosuppression. *The Plant Cell* 9: 1245-1249.

- Taylor, D.C., Weber, N., Barton, D.L., Underhill, E.W., Hogge, L.R., Weselake, R.J. and Pomeroy, M.K. (1991) Triacylglycerol bioassembly in microspore-derived embryos of *Brassica napus* L. cv. Reston. *Plant Physiol.* 97, 65-79.
- Taylor, D.C., Barton, D.L., Rioux, K.P., MacKenzie, S.L., Reed, D.W., Underhill, E.W., Pomeroy, M.K. and Weber, N. (1992) Biosynthesis of acyl lipids containing very-long chain fatty acids in microspore-derived embryos of *Brassica napus* L. cv. Reston. *Plant Physiol.* 99, 1609-1618.
- Tijburg, LB, Geelen, MJ and van Golde LM (1989) Regulation of the biosynthesis of triacylglycerol, phosphatidylcholine and phosphatidylethanolamine in the liver. *Biochim Biophys Acta* 1004: 1-19.
- Tzen TC, Cao Y, Laurent P, Ratnayake C and Huang HC (1993) Lipids, proteins and structures of seed oil bodies from diverse species. *Plant Physiol.* 101: 267-276.
- Vasil, I.K. (1994) Molecular improvement of cereals. *Plant Mol. Biol.* 25: 925-937.
- Vaucheret, H., Béclin C, Elmayan T, Feuerbach, F., Godon C, Morel J-B, Mourrain, P., Palauqui, J-C and Vernhettes S (1998) Transgene-induced gene silencing in plants. *The Plant Journal* 16: 651-659.
- Voelker, T.A., Worrell, A.C., Anderson, L., Bleibaum, J., Fan, C., Hawkins, D.J., Radke, S.E., and Davies, H.M. (1992) Fatty acid biosynthesis redirected to medium chains in transgenic oilseed plants. *Science* 257: 72-74.
- Voelker, T.A., Hayes, T.R., Cramner, A.M., Turner, J.C., and Davies, H.M. (1996) Genetic engineering of a quantitative trait: metabolic and genetic parameters influencing the accumulation of laurate in rapeseed. *The Plant Journal* 9: 229-241.
- Vogel, G and Browse, J (1996) Cholinephosphotransferase and diacylglycerol acyltransferase: Substrate specificities at a key branch point in seed lipid metabolism. *Plant Physiol.* 110: 923-931.
- Walden, R. and Wingender, R. (1995) Gene-transfer and plant regeneration techniques. *Trends in Biotechnology* 13: 324-331.

- Weselake, R.J., Taylor, D.C., Pomeroy, M.K., Lawson S.L. and Underhill E.W. (1991) properties of diacylglycerol acyltransferase from microspore-derived embryos of *Brassica napus* L. *Phytochemistry* : 30: 3533-3538.
- Weselake, R.J., Pomeroy, M.K., Furukawa, T.L., Golden, J.L., Little, D.B. and Laroche, A. (1993) Developmental profile of diacylglycerol acyltransferase in maturing seeds of oilseed rape and safflower and micro-spore-derived cultures of oilseed rape. *Plant Physiol.* 102, 565-571.
- Wilson, R.F. and Kwan-yuan P. (1986) Triacylglycerol synthesis and metabolism in germinating soybean cotyledons. *Biochim. Biophys. Acta* 877, 231-237.
- Yang, H., Bard, M., Bruner, D.A., Gleeson, A., Deckelbaum, R.J., Aljinovic, G., Pohl, T.M., Rothstein, R. and Sturley, S.L. (1997) Sterol esterification in yeast: a two-gene process. *Science* 272, 1353-1356.
- Yu, C., Kennedy, N.J., Chang, C.C.Y. and Rothblatt, J.A. (1996) Molecular cloning and characterization of two isoforms of *Saccharomyces cerevisiae* Acyl-CoA: Sterol Acyltransferase. *J. Biol. Chem.* 271, 24157-24163.
- Zou, J-T., Katavic, V., Giblin, E.M., Barton, D.L., MacKenzie, S.L., Keller, W.A., Hu, X. and Taylor, D.C. (1997) Modification of seed oil content and acyl composition in the Brassicaceae by expression of a yeast sn-2 acyltransferase gene. *The Plant Cell* 9: 909-923.

CLAIMS:

1. Isolated and purified deoxyribonucleic acid (DNA), characterized in that said DNA includes a sequence according to SEQ ID NO:1 or a part of SEQ ID NO:1, or a sequence that is substantially homologous to SEQ ID NO:1.
- 5 2. Isolated and purified deoxyribonucleic acid (DNA), characterized in that said DNA includes a sequence according to SEQ ID NO:3 or a part of SEQ ID NO:3, or a sequence that is substantially homologous to SEQ ID NO:3.
3. A vector that contains a nucleic acid sequence according to SEQ ID NO:1, or a part of SEQ ID NO:1, or a sequence that is substantially homologous to
10 SEQ ID NO:1.
4. A vector that contains a nucleic acid sequence according to SEQ ID NO:3, or a part of SEQ ID NO:3, or a sequence that is substantially homologous to SEQ ID NO:3.
5. A vector that contains a deoxyribonucleic acid sequence according to
15 SEQ ID NO:23.
6. A vector according to claim 3 or claim 5, characterized in that said sequence is present in said vector in a sense orientation.
7. A vector according to claim 3, characterized in that said sequence is present in said vector in an anti-sense orientation.
- 20 8. Plasmid pDGATcDNA (ATCC PTA-989).
9. Plasmid pDGATgene (ATCC PTA-988).
10. A plant having a genome, characterized in that the genome contains an introduced recombinant nucleotide sequence of SEQ ID NO:1, or a part of SEQ ID NO:1, or a sequence that is substantially homologous to SEQ ID NO:1.
- 25 11. A plant seed having a genome, characterized in that said genome contains an introduced recombinant nucleotide sequence of SEQ ID NO:1, or a part of SEQ ID NO:1, or a sequence that is substantially homologous to SEQ ID NO:1.

12. A genetically transformed plant, characterized in that said plant has a genome that has been transformed by a vector according to claim 3 or claim 4 or claim 5.

13. A genetically transformed plant seed, characterized in that said seed has been transformed by a vector according to claim 3 or claim 4 or claim 5.

14. A plant seed as claimed in Claim 11 or Claim 13, characterized by exhibiting an altered seed oil content compared to the seed oil content of seeds of genomically-unmodified plants of the same genotype grown in identical conditions at the same time.

10 15. A plant seed as claimed in Claim 11 or Claim 13, characterized by exhibiting an altered diacylglycerol content in its seed oil compared to the diacylglycerol content of seeds of genomically-unmodified plants of the same genotype grown in identical conditions at the same time.

15 16. A plant seed as claimed in 11 or 13, characterized by exhibiting a seed oil with an altered fatty acyl composition compared to the fatty acid composition of plants of the same genotype grown in identical conditions at the same time.

17. A plant as claimed in 10 or 12, characterized by exhibiting an enhanced biomass compared to the biomass of genomically-unmodified plants of the same genotype grown in identical conditions at the same time.

20 18. A seed as claimed in 11 or 13, characterized by exhibiting an enhanced biomass compared to the biomass of genomically-unmodified plants of the same genotype grown under identical conditions at the same time.

19. A method of producing transgenic plants by introducing a recombinant nucleotide sequence into a genome of said plant, characterized in that said nucleotide sequence introduced into said genome includes SEQ ID NO:1, SEQ ID NO:3 or SEQ ID NO:23; or a part of SEQ ID NO:1, SEQ ID NO:3 or SEQ ID NO:23; or a sequence that is substantially homologous to SEQ ID NO:1, SEQ ID NO:3 or SEQ ID NO:23.

20. A method according to claim 19, characterized in that said plant is a member of the Brassicaceae.

21. A method according to claim 19, characterized in that said plant is selected from the group consisting of *Arabidopsis thaliana*, borage (*Borago* spp.), Canola, castor (*Ricinus communis*), cocoa bean (*Theobroma cacao*), corn (*Zea mays*), cotton (*Gossypium* spp.), *Crambe* spp., *Cuphea* spp., flax (*Linum* spp.), *Lesquerella* 5 and *Limnanthes* spp., Linola, nasturtium (*Tropaeolum* spp.), *Oenothera* spp., olive (*Olea* spp.), palm (*Elaeis* spp.), peanut (*Arachis* spp.), rapeseed, safflower (*Carthamus* spp.), soybean (*Glycine* and *Soja* spp.), sunflower (*Helianthus* spp.), tobacco (*Nicotiana* spp.), *Vernonia* spp., wheat (*Triticum* spp.), barley (*Hordeum* spp.), rice (*Oryza* spp.), oat (*Avena* spp.) sorghum (*Sorghum* spp.), rye (*Secale* spp.) and other 10 members of the *Gramineae*.

22. An isolated plant DNA sequence or part thereof, characterized in that the sequence is substantially homologous to at least a part of SEQ ID NO:1 or SEQ ID NO:3.

15 23. A method of changing the oil content, acyl composition or diacylglycerol/triacylglycerol proportions of the seed oil of plant seeds by introducing a sense or anti-sense recombinant nucleic acid construct into a plant transformation vector, using the vector to transform the genome of a plant or plant seed, and then growing the plant or plant seed and extracting the oil from the plant seed, characterized in that said 20 recombinant nucleic acid sequence comprises SEQ ID NO:1, SEQ ID NO:3 or SEQ ID NO:23, or a part of SEQ ID NO:1, SEQ ID NO:3 or SEQ ID NO:23; or a sequence that is substantially homologous to SEQ ID NO:1, SEQ ID NO:3 or SEQ ID NO:23.

1/10




FIG. 1

2/10




FIG. 2




FIG. 3

3/10




FIG. 4

4/10

(a)

AtTAG1 MAILDSAGVTTVTENGGEFVDLDRLRRRKSRSDSSNGLLLSGSDNNSPS 50
MDGAT MEDRGGAGSSR.....RRRTGSRVSVQG....GSGPKVEE 31
HARGP1 MGDR...GSSR.....RRRTGSRPSSHG....GGGPAAAE 28

AtTAG1 DEVGAPADVDRIDSVVNDDAQGTANLAGDNNGGGDNNGGGRGCGGEGRGN 100
MDGAT DEVRDAAVSPDL.....GAGGDAPAPAPAPAHTRDKDGRTSVGEGY.. 72
HARGP1 DEVRDAAGPDV.....GAAGDAPAPAP.....NKDGDAGVGSCH.. 63

AtTAG1 ADATETYPSPVPAHRRARESPSSDAIFKQSHAGIFNL CVV LIAVNR L 150
MDGATWDLRC.....HRLQDSLFSDDGFSNYR.GILNWCVV LILSN R L 112
HARGP1WELRC.....HRLQDSLFSDDGFSNYR.GILNWCVV LILSN R L 103

AtTAG1 I IENLKYGWLIRTDWFSSRSRLRD...WPLFMC.CISLSIFPLAAFTVE 196
MDGAT FIENLKYGILVDP.IQVVSFLFKDPYSWPA.PCVIIASNIFVVAAFQIE 160
HARGP1 FIENLKYGILVDP.IQVVSFLFKDPHSWPA.PCLVIAANVEAFAAFQVE 151

AtTAG1 KLV LQKYI SEP VVI FLHI I ITMTEVLYPVYVTLRCD SAFLSGVTMLLTC 246
MDGAT KRIAVGALTEOMGLLLHVVNLAITICEPAAVALLVESITPVGVSFALASY 210
HARGP1 KRLAVGALTEOAGLLLVANLAITICEPAAVLLVESITPVGSLALMAH 201

AtTAG1 .IIVWLKLVSYAHTSY.....DIRSLANAADK.....ANPEVSYVYSL.. 282
MDGAT SIMLKLKLYSYRDVNLWCRRRVKAKAVSTGKKVSGAAAQQAVSYPDNLTY 260
HARGP1 TILKLKLYSYRDVNSWC..RRARAKAASAGKKASSAAAPHTVSYPDNLTY 249

AtTAG1 KSLAYFMVAPTLCYQPSYPRSA CIRKGWVARQFAKLVIFTFGMGFIEBOY 332
MDGAT RDLYYFIFAPTLCYELNEPRSPRIKRELLRRVLEMEFFTQLQVGLIQOW 310
HARGP1 RDLYYFIFAPTLCYELNEPRSPRIKRELLRILEMEFFTQLQVGLIQOW 299

AtTAG1 INPIVRNSKHPLKGDILYA..IERVLKLSVPNLYVWLCMFYCFHFLWLNI 380
MDGAT MVPTTONSMKPFFK.DMDYSRIIERELKLAVPNHLIWLIFFYWFFHSCLNA 359
HARGP1 MVPTTONSMKPFFK.DMDYSRIIERELKLAVPNHLIWLIFFYWLFHSCLNA 348

AtTAG1 LAELFCFGDREFYKDWNNAKSVGDYWRMWNMPVHKWVRHIYFPCLRSKI 430
MDGAT VAELFCFGDREFYRDWNNAESVTYEWQNNMPVHKWCTRHFYKPMRLRHGS 409
HARGP1 VAELFCFGDREFYRDWNNSESVTYEWQNNMPVHKWCTRHFYKPMRLRRGS 398

AtTAG1 PKTIAIIIAFLVSAVFHELCTAVPCRIFKLWAF LGIMFOVPLV.FETNYL 479
MDGAT SKWVARTGVFLTSAFFHEYLVSVP LRMFRLWAFTAMMAQVPLAWIVGREF 459
HARGP1 SKWMARTGVFLASAFFHEYLVSVP LRMFRLWAFTGMMAQIPLAWFVGRF 448

AtTAG1 QERFGSTVGNMIFWFIFCIFGQPMCVLIIYHD..LMNRKGSMS.. 520
MDGAT QGNYGNAA....VWVTL.IIGQPIAVLMYVHDYVVENYDAPVG.V 498
HARGP1 QGNYGNAA....VWLSL.IIGQPIAVLMYVHDYVVENYEAPAAEA 488

FIG. 5(a)




FIG. 5(b)


6/10

RT LF FL YS SL GS



FIG. 6(a)

1 2 3 bp



— 506

FIG. 6(b)

7/10




FIG. 7(a)




FIG. 7(b)

8/10




FIG. 8




FIG. 9

9/10




FIG. 10




FIG. 11

10/10




FIG. 12




FIG. 13

Please type a plus sign (+) inside this box → ☐

PTO/SB/01 (12-97)
Approved for use through 9/30/00. OMB 0651-0032
Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

**DECLARATION FOR UTILITY OR
DESIGN
PATENT APPLICATION
(37 CFR 1.63)**

☐ Declaration Submitted with Initial Filing OR ☒ Declaration Submitted after Initial Filing (surcharge (37 CFR 1.16 (e)) required)

Attorney Docket Number 43922
First Named Inventor ZOU, Jitao et al.

COMPLETE IF KNOWN

Application Number 09 / 623,514 /
Filing Date
Group Art Unit
Examiner Name

As a below named inventor, I hereby declare that:

My residence, post office address, and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

DIACYLGLYCEROL ACYLTRANSFERASE GENE FROM PLANTS

the specification of which (Title of the Invention)

☐ is attached hereto
OR

☒ was filed on (MM/DD/YYYY) 12/16/1999 as United States Application Number or PCT International

Application Number PCT/CA99/01202 and was amended on (MM/DD/YYYY) (if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment specifically referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR 1.56.

I hereby claim foreign priority benefits under 35 U.S.C. 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application Number(s)	Country	Foreign Filing Date (MM/DD/YYYY)	Priority Not Claimed	Certified Copy Attached?	
				YES	NO
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

☐ Additional foreign application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto:

I hereby claim the benefit under 35 U.S.C. 119(e) of any United States provisional application(s) listed below.

Application Number(s)	Filing Date (MM/DD/YYYY)	<input type="checkbox"/> Additional provisional application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto.
60/112,812 ✓	12/17/1998 ✓	

[Page 1 of 2]

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.

Please type a plus sign (+) inside this box → ☒

PTO/SB/01 (12-97)
Approved for use through 9/30/00. OMB 0651-0032
Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

DECLARATION — Utility or Design Patent Application

I hereby claim the benefit under 35 U.S.C. 120 of any United States application(s), or 365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. 112, I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

U.S. Parent Application or PCT Parent Number	Parent Filing Date (MM/DD/YYYY)	Parent Patent Number (if applicable)

☐ Additional U.S. or PCT international application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto.

As a named inventor, I hereby appoint the following registered practitioner(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

☐ Customer Number

OR

☒ Registered practitioner(s) name/registration number listed below

Place Customer
Number Bar Code
Label here

Name	Registration Number	Name	Registration Number
Edwin J. Gale	28,584	Kimberley A. Lachaine	33,319
Norris M. Eades	25,263	Robert K. Feutlinske	37,994
John A. Baker	26,656	Andrew J. Bauer-Moore	44,449

☐ Additional registered practitioner(s) named on supplemental Registered Practitioner Information sheet PTO/SB/02C attached hereto.

Direct all correspondence to: ☐ Customer Number or Bar Code Label

OR ☒ Correspondence address below

Name	EDWIN J. GALE,		
Address	KIRBY, EADES, GALE, BAKER,		
Address	P.O. Box 3432, Station D,		
City	Ottawa,	State	Ontario
Country	CANADA	Telephone	(613) 237-6900
		Fax	(613) 237-0045

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. 1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Name of Sole or First Inventor:

☒ A petition has been filed for this unsigned inventor

Given Name (first and middle (if any))	Family Name or Surname
JITAO	ZOU

Inventor's Signature	J. L. Zou		Date	Oct 18, 2000			
Residence: City	Saskatoon	State	SK	Country	CANADA	Citizenship	CAN.
Post Office Address	1619 Acadia Dr.						
Post Office Address							
City	Saskatoon	State	SK	ZIP	S7H 4T8	Country	CANADA

☒ Additional inventors are being named on the supplemental Additional Inventor(s) sheet(s) PTO/SB/02A attached hereto




Please type a plus sign (+) inside this box → +

PTO/SB/02A (3-97)
Approved for use through 9/30/98. OMB 0651-0032
Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

DECLARATION

ADDITIONAL INVENTOR(S) Supplemental Sheet Page 2 of 2

Name of Additional Joint Inventor, if any:				<input checked="" type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name (first and middle (if any))				Family Name or Surname			
2-00 <u>DAVID C.</u>				<u>TAYLOR</u>			
Inventor's Signature				Date	Oct 18, 2000		
Residence: City	<u>Saskatoon</u> <u>CAX</u>	State	SK	Country	CANADA	Citizenship	Canada ✓
Post Office Address	622 Wollaston Bay						
Post Office Address							
City	Saskatoon	State	SK	ZIP	S7J 4C3	Country	CANADA
Name of Additional Joint Inventor, if any:				<input checked="" type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name (first and middle (if any))				Family Name or Surname			
3-00 <u>YANGDOU</u>				<u>WEI</u>			
Inventor's Signature				Date	Oct 17, 2000		
Residence: City	<u>Saskatoon</u> <u>CAX</u>	State	SK	Country	CANADA	Citizenship	CAN. ✓
Post Office Address	1514 Main Street, Apt. 8						
Post Office Address							
City	Saskatoon	State	SK	ZIP	S7H 0L7	Country	Canada
Name of Additional Joint Inventor, if any:				<input checked="" type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name (first and middle (if any))				Family Name or Surname			
4-00 <u>COLETTE C.</u>				<u>JAKO</u>			
Inventor's Signature				Date	Oct 17, 2000		
Residence: City	<u>Saskatoon</u> <u>CAX</u>	State	SK	Country	CANADA	Citizenship	France <u>CAN.</u> <u>CI</u>
Post Office Address	1419 7th Avenue N.						
Post Office Address							
City	Saskatoon	State	SK	ZIP	S7K 2W3	Country	Canada

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.

SEQUENCE LISTING

<110> National Research Council of Canada
Zou, Jitao
Taylor, David C
Wei, Yangdou
Jako, Colette C

<120> Diacylglycerol Acyl Transferase Gene from Plants

<130> 43922
<140> US 09/623,514
<141> 2000-10-03
<150> PCT/CA99/01202
<151> 1999-12-16
<150> US 60/112,812
<151> 1998-12-17

<160> 25

<170> PatentIn Ver. 2.1

<210> 1
<211> 1904
<212> DNA
<213> Arabidopsis thaliana

<400> 1
atttcttagc ttcttccttc aatccgctct ttcctctctc attagattct gtttctcttt 60
tcaattttctt ctgcatgctt ctcgattctc tctgacgcct cttttctccc gacgctgttt 120
cgtcaaacgc ttttcgaaat ggcgattttg gattctgctg gcgttactac ggtgacggag 180
aacggtggcg gagagttcgt cgatcttgat aggtctcgtc gacggaaatc gagatcggat 240
tcttctaacg gacttcttct ctctggttcc gataataatt ctcttcgga tgatgttggg 300
gtccccgcgc acgttaggga tcggattgat tccgttgta acgatgacgc tcagggaaca 360
gccaatattg ccggagataa taacggtggt ggcgataata acggtggtgg aagaggcggc 420
ggagaaggaa gaggaacgc cgatgctacg tttacgtatc gaccgtcggt tccagctcat 480
cggagggcga gagagagtc acttagctcc gacgcaatct tcaaacagag ccatgccgga 540
ttattcaacc tctgtgtagt agttcttatt gctgtaaaca gtagactcat catcgaaaat 600
cttatgaagt atggttggtt gatcagaacg gatttctggt ttagttcaag atcgctgcga 660
gattggccgc ttttcatgtg ttgtatatcc ctttcgatct ttctttggc tgcctttacg 720
gttgagaaat tggtaactca gaaatacata tcagaacctg ttgtcatctt tcttcatatt 780
attatcacca tgacagaggt tttgtatcca gtttacgtca ccctaagggtg tgattctgct 840
tttttatcag gtgtcacttt gatgctcttc acttgcatg tgtggctaaa gttggtttct 900
tatgctcata ctagctatga cataagatcc ctagccaatg cagctgataa ggccaatcct 960
gaagtctcct actacgttag cttgaagagc ttggcatatt tcatggctgc tcccacattg 1020
tggtatcagc caagttatcc acgttctgca tgtatacgga agggttgggt ggctcgtcaa 1080
tttgcaaaac tggcatatt caccggattc atgggattta taatagaaca atatataaat 1140
cctattgtca ggaactcaaa gcatcctttg aaaggcgatc ttctatatgc tattgaaaga 1200
gtgttgaaagc tttcagttcc aaatttatat gtgtggctct gcatgttcta ctgcttcttc 1260
cacctttggt taaacatatt ggcagagctt ctctgcttcg gggatcgtga attctacaaa 1320
gattggtgga atgcaaaaag tgtgggagat tactggagaa tgtggaatat gcctgttcat 1380
aaatggatgg ttgcacatat atacttcccg tgcctgcgca gcaagatacc aaagacactc 1440
gccattatca ttgctttcct agtctctgca gtctttcatg agctatgcat cgcagttcct 1500
tgtcgtctct tcaagctatg ggccttttct gggattatgt ttcaggtgcc tttggtcttc 1560
atcacaaact atctacagga aagggttggt tcaacgggtg ggaacatgat cttctggttc 1620
atcttctgca ttttcgggaa accgatgtgt gtgcttcttt attaccacga cctgatgaac 1680
cgaaaaggat cgatgtcatg aaacaactgt tcaaaaaaat actttcttca aacatctatg 1740
gcctcggttg atctccgttg atgttgtggt ggttctgatg ctaaaacgac aaatagtgtt 1800
ataaccattg aagaagaaaa gaaaattaga gttgttgtat ctgcaaaaat tttggttagag 1860

1904

<211> 520

<212> PRT

<213> Arabidopsis thaliana

<400> 2

Met Ala Ile Leu Asp Ser Ala Gly Val Thr Thr Val Thr Glu Asn Gly
1 5 10 15

Gly Gly Glu Phe Val Asp Leu Asp Arg Leu Arg Arg Arg Lys Ser Arg
20 25 30

Ser Asp Ser Ser Asn Gly Leu Leu Leu Ser Gly Ser Asp Asn Asn Ser
35 40 45

Pro Ser Asp Asp Val Gly Ala Pro Ala Asp Val Arg Asp Arg Ile Asp
50 55 60

Ser Val Val Asn Asp Asp Ala Gln Gly Thr Ala Asn Leu Ala Gly Asp
65 70 75 80

Asn Asn Gly Gly Gly Asp Asn Asn Gly Gly Gly Arg Gly Gly Gly Glu
85 90 95

Gly Arg Gly Asn Ala Asp Ala Thr Phe Thr Tyr Arg Pro Ser Val Pro
100 105 110

Ala His Arg Arg Ala Arg Glu Ser Pro Leu Ser Ser Asp Ala Ile Phe
115 120 125

Lys Gln Ser His Ala Gly Leu Phe Asn Leu Cys Val Val Val Leu Ile
130 135 140

Ala Val Asn Ser Arg Leu Ile Ile Glu Asn Leu Met Lys Tyr Gly Trp
145 150 155 160

Leu Ile Arg Thr Asp Phe Trp Phe Ser Ser Arg Ser Leu Arg Asp Trp
165 170 175

Pro Leu Phe Met Cys Cys Ile Ser Leu Ser Ile Phe Pro Leu Ala Ala
180 185 190

Phe Thr Val Glu Lys Leu Val Leu Gln Lys Tyr Ile Ser Glu Pro Val
195 200 205

Val Ile Phe Leu His Ile Ile Ile Thr Met Thr Glu Val Leu Tyr Pro
210 215 220

Val Tyr Val Thr Leu Arg Cys Asp Ser Ala Phe Leu Ser Gly Val Thr
225 230 235 240

Leu Met Leu Leu Thr Cys Ile Val Trp Leu Lys Leu Val Ser Tyr Ala
245 250 255

His Thr Ser Tyr Asp Ile Arg Ser Leu Ala Asn Ala Ala Asp Lys Ala
260 265 270

Asn Pro Glu Val Ser Tyr Tyr Val Ser Leu Lys Ser Leu Ala Tyr Phe
 275 280 285

Met Val Ala Pro Thr Leu Cys Tyr Gln Pro Ser Tyr Pro Arg Ser Ala
 290 295 300

Cys Ile Arg Lys Gly Trp Val Ala Arg Gln Phe Ala Lys Leu Val Ile
 305 310 315 320

Phe Thr Gly Phe Met Gly Phe Ile Ile Glu Gln Tyr Ile Asn Pro Ile
 325 330 335

Val Arg Asn Ser Lys His Pro Leu Lys Gly Asp Leu Leu Tyr Ala Ile
 340 345 350

Glu Arg Val Leu Lys Leu Ser Val Pro Asn Leu Tyr Val Trp Leu Cys
 355 360 365

Met Phe Tyr Cys Phe Phe His Leu Trp Leu Asn Ile Leu Ala Glu Leu
 370 375 380

Leu Cys Phe Gly Asp Arg Glu Phe Tyr Lys Asp Trp Trp Asn Ala Lys
 385 390 395 400

Ser Val Gly Asp Tyr Trp Arg Met Trp Asn Met Pro Val His Lys Trp
 405 410 415

Met Val Arg His Ile Tyr Phe Pro Cys Leu Arg Ser Lys Ile Pro Lys
 420 425 430

Thr Leu Ala Ile Ile Ile Ala Phe Leu Val Ser Ala Val Phe His Glu
 435 440 445

Leu Cys Ile Ala Val Pro Cys Arg Leu Phe Lys Leu Trp Ala Phe Leu
 450 455 460

Gly Ile Met Phe Gln Val Pro Leu Val Phe Ile Thr Asn Tyr Leu Gln
 465 470 475 480

Glu Arg Phe Gly Ser Thr Val Gly Asn Met Ile Phe Trp Phe Ile Phe
 485 490 495

Cys Ile Phe Gly Gln Pro Met Cys Val Leu Leu Tyr Tyr His Asp Leu
 500 505 510

Met Asn Arg Lys Gly Ser Met Ser
 515 520

<210> 3

<211> 5193

<212> DNA

<213> Arabidopsis thaliana

<400> 3

gctcagcacc cattcttccc gttccatttg gttttattta tttcaaagtt taatattcct 60
 ttgtataac attcaaatct tcacatgatt gattgtgtga aaacccaca gattttacta 120

caataggggg	agttgactta	aaatagctat	tgatgtcgaa	aaaatgtatt	ttagttataa	180
attatactaa	agaaaatttt	tgatttgtct	gttgtttaag	catatgtatt	gttaaaactta	240
aaaaaatatg	tattgttaaat	cttaaaaatg	taggagtaca	catcaaatac	tcgagcataa	300
tcaaaaccgt	attcatagac	cgatgtgaga	atcaaataga	agataatgtg	atttttttaa	360
ataticgtatc	tccaaatcaa	tcacttagaa	gataatgtaa	ttctttatgt	gctacataaa	420
taaatatata	tatatatata	tatatatata	ttgtatatat	gtcttgacaa	aaaattgcc	480
gtcaaaaacc	atgactgaat	caaactataa	gtcggattga	atcaaactat	aagtcggatg	540
agtattaatt	tccattatgt	ttctatactt	tacaaaccgg	aaaatagata	ttatagatac	600
caaaaaagta	gatttgtgta	tattattaga	agatttggaa	tttcatcatt	atcaggatct	660
aaagtacttc	cctaattaaa	tcatgtcggg	tgaaaaagct	caatgaatgt	ttgaaatttg	720
gaaagtttat	taaattcgga	tctttttttt	ttgtttgtcg	tcccaaact	ttttatttta	780
ttacaaataa	tcaacttatc	cttactacta	aatcatttca	tatctttgat	accaacaaat	840
catttcatat	tctattttga	tgtttaagaa	aacactatct	accagttaca	aaatattata	900
aggattgttg	tttagaaaaa	aaagtacaag	ttgaattctt	tttgtcaa	ataaaattga	960
cttttttaata	tataattgac	ttattgaaca	tgattacaga	attaatcatc	tacaaaactt	1020
tccaaagtta	taataaatac	atttcaaaga	ctattagttc	ttcttaaaat	atttctaaaa	1080
gtgatcaaaag	actaccacat	ataattcaga	aaaagtagaa	gttgatttct	ttttgtcaaa	1140
taaataattg	acttaaaata	gtttggaaag	ccattgaact	tgattataga	attgataatg	1200
tacataaaaa	aattccaagt	ttataataaa	tacatttttc	aaatgctata	tcagttcttc	1260
ttaaaaatatt	tcactaaaaa	aacactcaaa	tatagaataa	atttattgaa	taacatacca	1320
actgtaaaaac	agaatttgac	aaaaaaaaaa	aaaaaatgaa	atgaagatga	agacaaaaat	1380
aaatcaccag	aggatcttat	gcaaaaaaat	atatgaatac	acaataaacc	atattgatat	1440
ttttaaaata	aaataaaaaac	agaaaaatat	cccaacaccg	cttttcaatt	aaaaatcttc	1500
cgtcaccatt	gttgtcatct	tctctctctg	tgaatccttt	ttcctttctt	cttcttcttc	1560
tcttcagaga	aaactttgct	tctctttcta	taaggaacca	gacacgaatc	ccattcccac	1620
cgattttotta	gcttcttctt	tcaatccgct	ctttccctct	ccattagatt	ctgtttcttc	1680
tttcaatttc	ttctgcatgc	ttctcgattc	tctctgacgc	ctcttttctc	ccgacgctgt	1740
ttcgtcaaac	gcttttcgaa	atggcgattt	tggattctgc	tggcgttact	acgggtgacg	1800
agaacggttg	cggagagttc	gtcgatcttg	ataggcttcg	tcgacggaaa	tcgagatcgg	1860
attcttctaa	cggacttctt	ctctctggtt	ccgataataa	ttctccttcg	gatgatgttg	1920
gagctcccg	cgacgttagg	gatcggattg	attccgttgt	taacgatgac	gctcagggaa	1980
cagccaattt	ggccggagat	aataacggtg	gtggcgataa	taacggtggt	ggaagaggcg	2040
gcgagagaagg	aagaggaaac	gccgatgcta	cgtttacgta	tcgacgctcg	gttccagctc	2100
atcggagggc	gagagagagt	ccacttagct	ccgacgcaat	cttcaaacag	gtttaaaatc	2160
tcagaaatct	tcgaatttgg	tgtttgcttg	ttgttttata	tgggaattgag	tttgggtgatt	2220
gttttgcat	gcagagccat	gccggattat	tcaacctctg	tgtagtagtt	cttattgctg	2280
taaacagtag	actcatcatc	gaaaatctta	tgaaggtttg	ctgttacttg	tttctccttt	2340
taggaattga	attgcttgaa	aatttatcag	agacgaataa	ctttgttgtt	gctatcatct	2400
atgtagtatg	gttggttgat	cagaacggat	ttctggttta	gttcaagatc	gctgcagatg	2460
tggcgcgttt	tcatgtgttg	gtaaaagaag	atgtttttta	ttccagcaa	tgtttacttg	2520
ttatacgtat	aatgatgagt	ttagtgtatc	agttcctctt	tgattcttct	ttcttgttgc	2580
agtatatccc	tttcgatctt	tcttttggtt	gcctttacgg	ttgagaaatt	gggtacttcag	2640
aaatacatat	cagaacctgt	gagtaattac	tattctccag	ccattactgt	aatttttatt	2700
gaagacaagt	ttgtatcatg	aagaacttac	aagttctggt	ttgaaaatgc	tcaaggttgt	2760
catctttctt	catattatta	tcaccatgac	agagggtttg	tatccagttt	acgtcaccct	2820
aagggtgatac	tgtttttctg	gtctcagttt	gtgatactgt	ttttaagttt	agttgtctga	2880
cccgggtgac	ttgaaaatgg	acagggtgtg	ttctgctttt	ttatcagggtg	tcactttgat	2940
gctcctcact	tgcattgtgt	ggctaaagtt	ggtttcttat	gctcatacta	gctatgacat	3000
aagatcccta	gccaatgcag	ctgataaggt	aaaatacgaa	aaagaagcgt	atgtattagt	3060
cacttgcact	gtgttactgt	tttaacccaa	cactgttatg	aacttttaggc	caatcctgaa	3120
gtctcctact	acgttagctt	gaagagcttg	gcataattca	tggctcgtcc	cacatttgtgt	3180
tatcaggtaa	ctgcaaagtg	catcaaccat	tcttatactt	gcaagagttt	cttgtctaaa	3240
cctcggatct	ttgcttttcc	ccagccaagt	tatccacggt	ctgcatgtat	acggaagggt	3300
tgggtggctc	gtcaatttgc	aaaactgggc	atattcaccg	gattcatggg	atttataata	3360
gaacaagtag	gttttcacat	cttgctttat	tagttttcct	tgggtgaaaat	catcatccct	3420
gcgttgctac	cacttgactt	catgttcttt	tgttactatt	tggcagttata	taaactctat	3480
tgtcagggaac	tcaaaagcatc	ctttgaaagg	cgatcttcta	tatgctattg	aaagagtgtt	3540
gaagctttca	gttccaaatt	tatatgtgtg	gctctgcatg	ttctactgct	tcttccacct	3600
ttggtatgct	gtgatcccat	ctctttcaaa	ataatttgca	aattcgaaaa	accgaaaaag	3660

Val Pro Leu Ala Trp Ile Val Gly Arg Phe Phe Gln Gly Asn Tyr Gly
450 455 460

Asn Ala Ala Val Trp Val Thr Leu Ile Ile Gly Gln Pro Val Ala Val
465 470 475 480

Leu Met Tyr Val His Asp Tyr Tyr Val Leu Asn Tyr Asp Ala Pro Val
485 490 495

Gly Val

<210> 5

<211> 488

<212> PRT

<213> human

<400> 5

Met Gly Asp Arg Gly Ser Ser Arg Arg Arg Arg Thr Gly Ser Arg Pro
1 5 10 15

Ser Ser His Gly Gly Gly Gly Pro Ala Ala Ala Glu Glu Glu Val Arg
20 25 30

Asp Ala Ala Ala Gly Pro Asp Val Gly Ala Ala Gly Asp Ala Pro Ala
35 40 45

Pro Ala Pro Asn Lys Asp Gly Asp Ala Gly Val Gly Ser Gly His Trp
50 55 60

Glu Leu Arg Cys His Arg Leu Gln Asp Ser Leu Phe Ser Ser Asp Ser
65 70 75 80

Gly Phe Ser Asn Tyr Arg Gly Ile Leu Asn Trp Cys Val Val Met Leu
85 90 95

Ile Leu Ser Asn Ala Arg Leu Phe Leu Glu Asn Leu Ile Lys Tyr Gly
100 105 110

Ile Leu Val Asp Pro Ile Gln Val Val Ser Leu Phe Leu Lys Asp Pro
115 120 125

His Ser Trp Pro Ala Pro Cys Leu Val Ile Ala Ala Asn Val Phe Ala
130 135 140

Val Ala Ala Phe Gln Val Glu Lys Arg Leu Ala Val Gly Ala Leu Thr
145 150 155 160

Glu Gln Ala Gly Leu Leu Leu His Val Ala Asn Leu Ala Thr Ile Leu
165 170 175

Cys Phe Pro Ala Ala Val Val Leu Leu Val Glu Ser Ile Thr Pro Val
180 185 190

Gly Ser Leu Leu Ala Leu Met Ala His Thr Ile Leu Phe Leu Lys Leu
195 200 205

092544 "0000" 44552360

<222> (576)..(576)
 <223> n is any nucleotide

<220>
 <221> misc_feature
 <222> (583)..(584)
 <223> n is any nucleotide

<220>
 <221> misc_feature
 <222> (597)..(597)
 <223> n is any nucleotide

<400> 6
 tgcattgtata cggaagggtt ggggtggctcg tcaatttgca aaactgggtca tattcaccgg 60
 attcatggga ttataaatag aacaatatat aaatcctatt gtcaggaact caaagcatcc 120
 ttgaaaggc gatcttctat atgctattga aagagtgttg aagctttcag ttccaaattt 180
 atatgtgtgg ctctgcatgt tctactgctt cttccacctt tggttaaaca tattggcaga 240
 gcttctctgc ttcggggatc gtgaattcta caaagattgg tggaatgcaa aaagtgtggg 300
 agattactgg gagaatgtgg aatatgcctg tccataaatg ggatgggtcc gacatatata 360
 ccttccccgt gcttgcgcac aaggattacc caaagacacc ccggccatta accattggct 420
 ttccaagcc ccttgaggc ctttccatgg gccanggacc cggngtnccc tggcngggcc 480
 ttcaaagcaa agggggnttn cctggggnta aagntccang ggcccttggg gcccanccaa 540
 aannttcccc cgggaaaggg ttgcccaccg gggggngaaa aanncccggg ggcaccncgg 600
 aattttggga acccgggggg ggccttttt 629

<210> 7
 <211> 11
 <212> PRT
 <213> Arabidops thaliana

<400> 7
 His Lys Trp Met Val Arg His Ile Tyr Phe Pro
 1 5 10

<210> 8
 <211> 341
 <212> PRT
 <213> Brassica napus

<400> 8
 Met Cys Cys Leu Ser Leu Ser Ile Phe Pro Leu Ala Ala Phe Thr Val
 1 5 10 15
 Glu Lys Leu Val Leu Gln Lys Cys Ile Ser Glu Pro Val Val Ile Phe
 20 25 30

Leu His Val Ile Ile Thr Met Thr Glu Val Leu Tyr Pro Val Tyr Val
 35 40 45
 Thr Leu Arg Cys Asp Ser Ala Phe Leu Ser Gly Asp Thr Leu Met Leu
 50 55 60
 Leu Thr Cys Ile Val Trp Leu Lys Leu Val Ser Tyr Ala His Thr Asn
 65 70 75 80
 Tyr Asp Ile Arg Thr Leu Ala Asn Ser Ser Asp Lys Ala Asn Pro Glu
 85 90 95
 Val Ser Tyr Tyr Val Ser Leu Lys Ser Leu Ala Tyr Phe Met Leu Ala
 100 105 110
 Pro Thr Leu Cys Tyr Gln Pro Ser Tyr Pro Arg Ser Pro Cys Ile Arg
 115 120 125
 Lys Gly Trp Val Ala Arg Gln Phe Ala Lys Leu Val Ile Phe Thr Gly
 130 135 140
 Leu Met Gly Phe Ile Ile Glu Gln Tyr Ile Asn Pro Ile Val Arg Asn
 145 150 155 160
 Ser Lys His Pro Leu Lys Gly Asp Leu Leu Tyr Ala Ile Glu Arg Val
 165 170 175
 Leu Lys Leu Ser Val Pro Asn Leu Tyr Val Trp Leu Cys Met Phe Tyr
 180 185 190
 Cys Phe Phe His Leu Trp Leu Asn Ile Leu Ala Glu Leu Leu Cys Phe
 195 200 205
 Gly Asp Arg Glu Phe Tyr Lys Asp Trp Trp Asn Ala Lys Ser Val Gly
 210 215 220
 Asp Tyr Trp Arg Met Trp Asn Met Pro Val His Lys Trp Met Val Arg
 225 230 235 240
 His Val Tyr Phe Pro Cys Leu Arg Ile Lys Ile Pro Lys Val Pro Ala
 245 250 255
 Ile Ile Ile Ala Phe Leu Val Ser Ala Val Phe His Glu Leu Cys Ile
 260 265 270
 Ala Val Pro Cys Arg Leu Phe Asn Leu Trp Ala Phe Met Gly Ile Met
 275 280 285
 Phe Gln Val Pro Leu Val Phe Ile Thr Asn Phe Leu Gln Glu Arg Phe
 290 295 300
 Gly Ser Met Val Gly Asn Met Ile Phe Gly Ser Ala Ser Cys Ile Phe
 305 310 315 320
 Gly Gln Pro Met Cys Gly Leu Leu Tyr Tyr His Asp Leu Met Asn Arg
 325 330 335
 Lys Gly Ser Met Ser

"000000" 47522560


```
<210> 10
<211> 11
<212> PRT
<213> Arabidopsis thaliana

<220>
<221> misc_feature
<222> (1)..(1)
<223> Xaa is any amino acid
```


<220>
 <221> misc_feature
 <222> (3)..(3)
 <223> Xaa is any amino acid

<220>
 <221> misc_feature
 <222> (5)..(6)
 <223> Xaa is any amino acid

<220>
 <221> misc_feature
 <222> (8)..(10)
 <223> Xaa is any amino acid

<400> 10
 Xaa Leu Xaa Leu Xaa Xaa Ser Xaa Xaa Xaa Val
 1 5 10

<210> 11
 <211> 27
 <212> PRT
 <213> Arabidopsis thaliana

<400> 11
 Ser His Ala Gly Leu Phe Asn Leu Cys Val Val Val Leu Ile Ala Val
 1 5 10 15

Asn Ser Arg Leu Ile Ile Glu Asn Leu Met Lys
 20 25

<210> 12
 <211> 24
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer of
 DGATXbal

<400> 12
 ctagtctaga atggcgattt tgga

24

<210> 13
 <211> 23
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer of
 DGATXhol

<400> 13

gcgctcgagt ttcatgacat cga

23

<210> 14

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer DGAT1

<400> 14

agacacgaat cccattccca ccga

24

<210> 15

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer DGAT2

<400> 15

agtggtgaca acgcagggat gatg

24

<210> 16

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer DGAT3

<400> 16

atggtcgctc ccacattgtg t

21

<210> 17

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer DGAT4

<400> 17

catacaatcc ccatgacatt tatca

25

<210> 18

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer A

<400> 18
cgaccgtcgg ttccagctca tcgg 24

<210> 19
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer B

<400> 19
gcggccaatc tcgcagcgat cttg 24

<210> 20
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer C

<400> 20
taaacagtag actcatcatc g 21

<210> 21
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer Gen 1

<400> 21
gagaggatcc acgctcacga cccattcttc ccg 33

<210> 22
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer Gen 2

<400> 22
aagaaggatc catccccaaa acgggaccac caa 33

<210> 23
<211> 1985
<212> DNA
<213> Arabidopsis thaliana

<400> 23
atttcttagc ttcttccttc aatccgctct ttccctctcc attagattct gtttcctctt 60

tcaattttctt ctgcatgctt ctcgattctc tctgacgcct cttttctccc gacgctgttt 120
 cgtcaaaccgc ttttcgaaat ggcgattttg gattctgctg gcgttactac ggtgacggag 180
 aacggtggcg gagagttcgt cgatcttgat aggcttcgtc gacggaaatc gagatcggat 240
 tcttctaacc gacttcttct ctctgggtcc gataataatt ctcttcgga tgatgttgga 300
 gctcccgccg acgttaggga tcggattgat tccgttgta acgatgacgc tcagggaaca 360
 gccaatttgg ccggagataa taacgggtgg ggcgataata acggtggtgg aagaggcggc 420
 ggagaaggaa gaggaaccgc cgatgctacg tttacgtatc gaccgtcggg tccagctcat 480
 cggagggcgga gagagagtcc acttagctcc gacgcaatct tcaaacagag ccatgccgga 540
 ttattcaacc tctgtgtagt agttcttatt gctgtaaaca gtagactcat catcgaaaat 600
 cttatgaaga gccatgccgg attattcaac ctctgtgtag tagttcttat tgctgtaaac 660
 agtagactca tcatcgaaaa tcttatgaag tatggttggg tgatcagaac ggatttctgg 720
 tttagttcaa gatcgctgcg agattggcgg cttttcatgt gttgtatatc cctttcgatc 780
 tttcctttgg ctgcctttac ggttgagaaa ttggtacttc agaaatacat atcagaacct 840
 gttgtcatct ttcttcatat tattatcacc atgacagagg ttttgtatcc agtttacgtc 900
 accctaagggt gtgattctgc ttttttatca ggtgtcactt tgatgctcct cacttgcatt 960
 gtgtggctaa agttggtttc ttatgctcat actagctatg acataagatc cctagccaat 1020
 gcagctgata aggccaatcc tgaagtctcc tactacgtta gcttgaagag cttggcataat 1080
 ttcattggtcg ctcccacatt gtgttatcag ccaagttatc cacgttctgc atgtatacgg 1140
 aagggttggg tggtcgtca atttgcaaaa ctggtcatat tcaccggatt catgggattt 1200
 ataatagaac aatatataaa tctattgtc aggaactcaa agcatccttt gaaaggcgat 1260
 cttctatatg ctattgaaag agtggtgaag ctttcagttc caaatttata tgtgtggctc 1320
 tgcattgtct actgcttctt ccacttttgg ttaaaccat tggcagagct tctctgcttc 1380
 ggggatcgtg aattctacaa agattggtgg aatgcaaaaa gtgtgggaga ttactggaga 1440
 atgtggaata tgctgttca taaatggatg gttcgacata tatacttccc gtgcttgcgc 1500
 agcaagatac caaagacact cgccattatc attgctttcc tagtctctgc agtctttcat 1560
 gagctatgca tcgcagttcc ttgtcgtctc ttcaagctat gggcttttct tgggattatg 1620
 tttcaggtgc ctttgggtct catcacaaac tatctacagg aaagggttgg ctcaacgggtg 1680
 gggaacatga tcttctggtt catcttctgc attttcggac aaccgatgtg tgtgcttctt 1740
 tattaccacg acctgatgaa ccgaaaagga tcgatgtcat gaaacaactg ttcaaaaaat 1800
 gactttcttc aaacatctat ggcctcgttg gatctccgtt gatgttgtgg tggttctgat 1860
 gctaaaacga caaatagtgt tataaccatt gaagaagaaa agaaaattag agttgttgta 1920
 tctgcaaaaa ttttggtaga gacacgcaaa cccgtttgga ttttgttatg gtgtaaagcg 1980
 gccgc 1985

<210> 24

<211> 5339

<212> DNA

<213> *Arabidopsis thaliana*

<400> 24

gctcaccgacc cattcttccc gttccatttg gttttattta tttcaaagtt taatattcct 60
 tttgtataac attcaaactc tcacatgatt gattgtgtga aaaccccaca gattttacta 120
 caataggggg agttgactta aaatagctat tgatgtcgaa aaaatgtatt ttagttataa 180
 attatactaa agaaaaat tttt gattttgtct gttgtttaag catatgtatt gttaaactta 240
 aaaaaatag tattgttaat cttaaaaaat taggagtaca catcaaatac tcgagcataa 300
 tcaaaaccgt attcatagac cgatgtgaga atcaaataga agataatgtg attttttaaa 360
 atatcgatc tccaaatcaa tcacttagaa gataatgtaa ttctttatgt gctacataaa 420
 taaatatata tatatatata tatatatatc ttgtatatat gtcttgacaa aaaattgcca 480
 gtcaaaaacc atgactgaat caaactataa gtcggattga atcaaactat aagtcggatg 540
 agtattaatt tccattatgt ttctatactt tacaaaccgg aaaatagata ttatagatac 600
 caaaaaagta gatttgtgta tattattaga agatttggaa tttcatcatt atcaggatct 660
 aaagtacttc cctaattaaa tcatgtcggg tgaaaaagct caatgaatgt ttgaaatttg 720
 gaaagtttat taaattcggg tctttttttt ttgtttgtcg tcccaaacat ttttatttta 780
 ttacaaataa tcaacttatc cttactataa aatcatttca tatctttgat accaacaata 840
 catttcatat tctattttga tgttttaagaa aaactattt accagttaca aaatattata 900
 aggattgttg tttagaaaaa aaagtacaag ttgaattctt tttgtcaaat ataaaaattga 960
 ctttttaata tataattgac ttattgaaca tgattacaga attaatcatc tacaaaaactt 1020
 tccaagttta taataaatac atttcaaaga ctattagttc ttcttaaaat atttctaaaa 1080

ctgaccaacc aatttgtttg agtaggtgcc tttggtcttc atcacaaact atctacagga 4680
 aagggtttggc tcaacgggtat gctctcaaaa cccgagaaaa tagaacgaat aactctttct 4740
 ttcatagcct agccatttaa atcgcaatgc tgaacttaa taataaagggt gatctgtttt 4800
 ggaatgggat catattatta ggtggggaac atgatcttct ggttcatctt ctgcattttc 4860
 ggacaaccga tgtgtgtgct tctttattac cacgacctga tgaaccgaaa aggatcgatg 4920
 tcatgaaaca actgttcaaa aaatgacttt cttcaaacaat ctatggcctc gttggatctc 4980
 cgttgatgtt gtggtggttc tgatgctaaa acgacaaata gtgttataac cattgaagaa 5040
 gaaaagaaaa ttagagttgt tgtatctgca aaaattttgg tagagacacg cgaaccggtt 5100
 tggattttgt tatggtgtaa agaaatttca atcaaaaaaac tgttgtaata attgttacca 5160
 aaaagaaatg cttttctgga aacgagggga aaaatagtag ttttgtagg ttttactggt 5220
 tggaccaaatt ctagtataaaa actttttgta ataaggaaaa aaaaagaaca aatgtgataa 5280
 atgcatgggg attgtatgaa accttccaat aaagttgatt ggtggtcccg ttttgggga 5339

<210> 25

<211> 547

<212> PRT

<213> Arabidopsis thaliana

<400> 25

Met Ala Ile Leu Asp Ser Ala Gly Val Thr Thr Val Thr Glu Asn Gly
 1 5 10 15

Gly Gly Glu Phe Val Asp Leu Asp Arg Leu Arg Arg Arg Lys Ser Arg
 20 25 30

Ser Asp Ser Ser Asn Gly Leu Leu Ser Gly Ser Asp Asn Asn Ser
 35 40 45

Pro Ser Asp Asp Val Gly Ala Pro Ala Asp Val Arg Asp Arg Ile Asp
 50 55 60

Ser Val Val Asn Asp Asp Ala Gln Gly Thr Ala Asn Leu Ala Gly Asp
 65 70 75 80

Asn Asn Gly Gly Gly Asp Asn Asn Gly Gly Gly Arg Gly Gly Gly Glu
 85 90 95

Gly Arg Gly Asn Ala Asp Ala Thr Phe Thr Tyr Arg Pro Ser Val Pro
 100 105 110

Ala His Arg Arg Ala Arg Glu Ser Pro Leu Ser Ser Asp Ala Ile Phe
 115 120 125

Lys Gln Ser His Ala Gly Leu Phe Asn Leu Cys Val Val Val Leu Ile
 130 135 140

Ala Val Asn Ser Arg Leu Ile Ile Glu Asn Leu Met Lys Ser His Ala
 145 150 155 160

Gly Leu Phe Asn Leu Cys Val Val Val Leu Ile Ala Val Asn Ser Arg
 165 170 175

Leu Ile Ile Glu Asn Leu Met Lys Tyr Gly Trp Leu Ile Arg Thr Asp
 180 185 190

Phe Trp Phe Ser Ser Arg Ser Leu Arg Asp Trp Pro Leu Phe Met Cys
 195 200 205

"00000" 45555555

Pro Met Cys Val Leu Leu Tyr Tyr His Asp Leu Met Asn Arg Lys Gly
530 535 540

Ser Met Ser
545

105200 4752950

MARKED-UP COPY

1
SHOWING AMENDMENTS

SEQUENCE LISTING

<110> National Research Council of Canada
Zou, Jitao
Taylor, David C
Wei, Yangdou
Jako, Colette C

<120> Diacylglycerol Acyl Transferase Gene from Plants

<130> ~~43922~~ 43922

<140> PCT/CA US 09/623,514

<141> 1999-12-16 2000-10-03

<150> 60/112,812

<151> 1998-12-17

(150) PCT/CA99/01202
(151) 1999-12-16

<160> 25

<170> PatentIn Ver. 2.1

<210> 1

<211> 1904

<212> DNA

<213> Arabidopsis thaliana

<400> 1

atttcttagc	ttcttcttc	aatcgcgtct	ttccctctcc	attagattct	gtttctctct	60
tcaatttctt	ctgcatgctt	ctcgattctc	tctgaagcct	cttttctccc	gacgctgttt	120
cgtcaaacgc	ttttcgaaat	ggcgattttg	gattctgctg	gcgttactac	ggtgacggag	180
aacggtggcg	gagagttcgt	cgatcttgat	aggcttcgtc	gacggaaatc	gagatcggat	240
tcttctaacg	gacttcttct	ctctggttcc	gataataatt	ctccttcgga	tgatgttgga	300
gtccccgcgc	acgttaggga	tcggattgat	tccgttggtt	acgatgacgc	tcagggaaca	360
gccaaatttg	ccggagataa	taacggtggt	ggcgataata	acggtggtgg	aagaggcggc	420
ggagaaggaa	gaggaaacgc	cgatgctacg	tttacgtatc	gaccgtcggg	tccagctcat	480
cggagggcga	gagagagtc	acttagctcc	gacgcaatct	tcaaacagag	ccatgccgga	540
ttattcaacc	tctgtgtagt	agttcttatt	gctgtaaaca	gtagactcat	catcgaaaat	600
cttatgaagt	atggttggtt	gatcagaacg	gatttctggt	ttagttcaag	atcgctgoga	660
gattggccgc	ttttcatgtg	ttgtatatcc	ctttcgatct	ttcctttggc	tgccctttacg	720
gttgagaaat	tggtacttca	gaaatacata	tcagaacctg	ttgtcatctt	tcttcatatt	780
attatcacca	tgacagaggt	tttgtatcca	gtttacgtca	ccctaagggtg	tgattctgct	840
tttttatcag	gtgtcacttt	gatgctcctc	acttgcatgt	tgtggctaaa	gttggtttct	900
tatgctcata	ctagctatga	cataagatcc	ctagccaatg	cagctgataa	ggccaatcct	960
gaagtctcct	actacgttag	cttgaagagc	ttggcatatt	tcattggtcgc	tcccacattg	1020
tgttatcagc	caagttatcc	acgttctgca	tgtatacggg	aggggtgggt	ggctcgtcaa	1080
tttgcaaaac	tggtcatatt	caccggattc	atgggattta	taatagaaca	atatataaat	1140
cctattgtca	ggaactcaaa	gcattcctttg	aaaggcgatc	ttctatatgc	tattgaaaga	1200
gtgttggaagc	tttcagttcc	aaatttataat	gtgtggctct	gcattgttcta	ctgcttcttc	1260
cacctttggt	taaacaatatt	ggcagagctt	ctctgcttcg	gggatcgtga	attctacaaa	1320
gattggtgga	atgcaaaaag	tgtgggagat	tactggagaa	tgtggaatat	gcctgttcat	1380
aaatggatgg	ttcgacatat	atacttcccg	tgtttgcgca	gcaagatacc	aaagacactc	1440
gccattatca	ttgctttcct	agtctctgca	gtctttcatg	agctatgcat	cgcagttcct	1500
tgtcgtctct	tcaagctatg	ggcttttctt	gggattatgt	ttcagggtgcc	tttgggtctc	1560
atcacaaact	atctacagga	aagggtttggc	tcaacggtgg	ggaacatgat	cttctgggtc	1620
atcttctgca	ttttcggaca	accgatgtgt	gtgcttcttt	attaccacga	cctgatgaac	1680
cgaaaaggat	cgatgtcatg	aaacaactgt	tcaaaaaatg	actttcttca	aacatctatg	1740

092514.03901

gcctcgttgg atctccgttg atgttggtggt ggttctgatg ctaaaacgac aaatagtgtt 1800
 ataaccattg aagaagaaaa gaaaattaga gttgttgat ctgcaaaaat tttggttagag 1860
 acacgcaaac ccgtttggat tttgttatgg tgtaaagcgg ccgc 1904

<210> 2
 <211> 520
 <212> PRT
 <213> Arabidopsis thaliana

<400> 2
 Met Ala Ile Leu Asp Ser Ala Gly Val Thr Thr Val Thr Glu Asn Gly
 1 5 10 15
 Gly Gly Glu Phe Val Asp Leu Asp Arg Leu Arg Arg Arg Lys Ser Arg
 20 25 30
 Ser Asp Ser Ser Asn Gly Leu Leu Leu Ser Gly Ser Asp Asn Asn Ser
 35 40 45
 Pro Ser Asp Asp Val Gly Ala Pro Ala Asp Val Arg Asp Arg Ile Asp
 50 55 60
 Ser Val Val Asn Asp Asp Ala Gln Gly Thr Ala Asn Leu Ala Gly Asp
 65 70 75 80
 Asn Asn Gly Gly Gly Asp Asn Asn Gly Gly Gly Arg Gly Gly Gly Glu
 85 90 95
 Gly Arg Gly Asn Ala Asp Ala Thr Phe Thr Tyr Arg Pro Ser Val Pro
 100 105 110
 Ala His Arg Arg Ala Arg Glu Ser Pro Leu Ser Ser Asp Ala Ile Phe
 115 120 125
 Lys Gln Ser His Ala Gly Leu Phe Asn Leu Cys Val Val Val Leu Ile
 130 135 140
 Ala Val Asn Ser Arg Leu Ile Ile Glu Asn Leu Met Lys Tyr Gly Trp
 145 150 155 160
 Leu Ile Arg Thr Asp Phe Trp Phe Ser Ser Arg Ser Leu Arg Asp Trp
 165 170 175
 Pro Leu Phe Met Cys Cys Ile Ser Leu Ser Ile Phe Pro Leu Ala Ala
 180 185 190
 Phe Thr Val Glu Lys Leu Val Leu Gln Lys Tyr Ile Ser Glu Pro Val
 195 200 205
 Val Ile Phe Leu His Ile Ile Ile Thr Met Thr Glu Val Leu Tyr Pro
 210 215 220
 Val Tyr Val Thr Leu Arg Cys Asp Ser Ala Phe Leu Ser Gly Val Thr
 225 230 235 240
 Leu Met Leu Leu Thr Cys Ile Val Trp Leu Lys Leu Val Ser Tyr Ala

"0625" 4152550

245										250					255				
His	Thr	Ser	Tyr	Asp	Ile	Arg	Ser	Leu	Ala	Asn	Ala	Ala	Asp	Lys	Ala				
			260					265					270						
Asn	Pro	Glu	Val	Ser	Tyr	Tyr	Val	Ser	Leu	Lys	Ser	Leu	Ala	Tyr	Phe				
		275					280						285						
Met	Val	Ala	Pro	Thr	Leu	Cys	Tyr	Gln	Pro	Ser	Tyr	Pro	Arg	Ser	Ala				
		290				295					300								
Cys	Ile	Arg	Lys	Gly	Trp	Val	Ala	Arg	Gln	Phe	Ala	Lys	Leu	Val	Ile				
305					310					315					320				
Phe	Thr	Gly	Phe	Met	Gly	Phe	Ile	Ile	Glu	Gln	Tyr	Ile	Asn	Pro	Ile				
				325					330					335					
Val	Arg	Asn	Ser	Lys	His	Pro	Leu	Lys	Gly	Asp	Leu	Leu	Tyr	Ala	Ile				
			340					345					350						
Glu	Arg	Val	Leu	Lys	Leu	Ser	Val	Pro	Asn	Leu	Tyr	Val	Trp	Leu	Cys				
		355					360					365							
Met	Phe	Tyr	Cys	Phe	Phe	His	Leu	Trp	Leu	Asn	Ile	Leu	Ala	Glu	Leu				
	370					375					380								
Leu	Cys	Phe	Gly	Asp	Arg	Glu	Phe	Tyr	Lys	Asp	Trp	Trp	Asn	Ala	Lys				
385					390					395					400				
Ser	Val	Gly	Asp	Tyr	Trp	Arg	Met	Trp	Asn	Met	Pro	Val	His	Lys	Trp				
			405						410					415					
Met	Val	Arg	His	Ile	Tyr	Phe	Pro	Cys	Leu	Arg	Ser	Lys	Ile	Pro	Lys				
			420					425					430						
Thr	Leu	Ala	Ile	Ile	Ile	Ala	Phe	Leu	Val	Ser	Ala	Val	Phe	His	Glu				
		435					440					445							
Leu	Cys	Ile	Ala	Val	Pro	Cys	Arg	Leu	Phe	Lys	Leu	Trp	Ala	Phe	Leu				
		450				455					460								
Gly	Ile	Met	Phe	Gln	Val	Pro	Leu	Val	Phe	Ile	Thr	Asn	Tyr	Leu	Gln				
465					470					475					480				
Glu	Arg	Phe	Gly	Ser	Thr	Val	Gly	Asn	Met	Ile	Phe	Trp	Phe	Ile	Phe				
				485					490					495					
Cys	Ile	Phe	Gly	Gln	Pro	Met	Cys	Val	Leu	Leu	Tyr	Tyr	His	Asp	Leu				
			500					505					510						
Met	Asn	Arg	Lys	Gly	Ser	Met	Ser												
			515				520												

<210> 3

<211> 5193

09554-03901

<212> DNA

<213> Arabidopsis thaliana

<400> 3

```

gctcacgacc cattcttccc gttccatttg gttttattta tttcaaagtt taatattcct 60
tttgtataac attcaaactc tcacatgatt gatttgtgtga aaacccacac gattttacta 120
caataggggg agttgactta aaatagctat tgatgtcgaa aaaatgtatt ttagttataa 180
attatactaa agaaaatttt tgatttgtct gttgtttaag catatgtatt gttaaactta 240
aaaaaatatg tattgttaat cttaaaaatg taggagtaca catcaaatac tcgagcataa 300
tcaaaaccgt attcatagac cgatgtgaga atcaaataga agataatgtg atttttttaa 360
atatcgatc tccaaatcaa tcacttagaa gataatgtaa ttctttatgt gctacataaa 420
taaatatata tatatatata tatatatatc ttgtatatat gtcttgacaa aaaattgcc 480
gtcaaaaacc atgactgaat caaactataa gtcggattga atcaaactat aagtcggatg 540
agtattaatt tccattatgt ttctatactt tacaaaaccg aaaatagata ttatagatac 600
caaaaaagta gatttgtgta tattattaga agatttggaa tttcatcatt atcaggatct 660
aaagtacttc cctaattaaa tcatgtcggg tgaaaaagct caatgaatgt ttgaaatttg 720
gaaagtttat taaattcgga tctttttttt ttgtttgtcg tcccaaacat ttttatttta 780
ttacaaataa tcaacttatc ctactacta aatcatttca tatctttgat accaacaat 840
catttcatat tctatttga tgtttaagaa aacactattt accagttaca aaatattata 900
aggattgttg tttagaaaaa aaagtacaag ttgaattctt tttgtcaaat ataaaaattga 960
ctttttaata tataattgac ttattgaaca tgattacaga attaatcatc taaaaaactt 1020
tccaagttaa taataaatac atttcaaaga ctattagtct ttcttaaaat atttctaaaa 1080
gtgatcaaag actaccacat ataattcaga aaaagtagaa gttgatttct ttttgtcaaa 1140
taaataattg acttaaaata gtttggaag ccattgaact tgattataga attgataatg 1200
tacataaaaa aattccaagt ttataataaa tacatttttc aaatgctata tcagttcttc 1260
ttaaaatatt tcaactaaaa aacactcaaa tatagaataa atttattgaa taacatacca 1320
actgtaaaac agaatttgac aaaaaaaaaa aaaaaatgaa atgaagatga agacaaaaat 1380
aaatcaccag aggatcttat gcaaaaaaat atatgaatac acaataaacc atattgatat 1440
ttttaaaata aaataaaaaa agaaaaatat cccaacaccg cttttcaatt aaaaatcttc 1500
cgtcaccatt gttgtcatct tctctctcgc tgaatccttt ttctttctt cttcttcttc 1560
tcttcagaga aaactttgct tctctttcta taaggaaaca gacacgaatc ccattcccac 1620
cgattttctt gcttcttcct tcaatccgct ctttccctct ccattagatt ctgtttcctc 1680
tttcaatttc ttctgcatgc ttctcgatc tctctgacgc ctcttttctc ccgacgctgt 1740
ttcgtcaaac gcttttcgaa atggcgattt tggattctgc tggcgttact acggtgacgg 1800
agaacggttg cggagagttc gtcgatcttg ataggcttcg tcgacggaaa tcgagatcgg 1860
attcttctaa cggacttctt ctctctgggt ccgataataa ttctccttcg gatgatgttg 1920
gagctcccg cgacgttagg gatcggattg attccgttgt taacgatgac gctcagggaa 1980
cagccaattt ggccggagat aataacgggt gtggcgataa taacggtggt ggaagaggcg 2040
gcgagagaag aagaggaaac gccgatgcta cgtttacgta tcgaccgtcg gttccagctc 2100
atcggagggc gagagagagt ccacttagct ccgacgcaat cttcaaacag gtttaaaatc 2160
tcagaaatct tcgaatttg tgtttgcttg ttgttttata tggaaattgag tttggtgatt 2220
gttttgcat gcagagccat gccggattat tcaacctctg tgtagtagtt cttattgctg 2280
taaacagtag actcatcatc gaaaatctta tgaaggtttg ctgttacttg tttctccttt 2340
taggaattga attgcttgaa aatttatcag agacgaataa ctttgttgtt gctatcattc 2400
atgtagtatg gttggttgat cagaacggat ttctggttta gttcaagatc gctgcgagat 2460
tgcccgcttt tcatgtgttg gtaaaaagaag atgtttttta tttccagcaa tgttacattg 2520
ttatacgtat aatgatgagt ttagtgatca agttcctctt tgattcttct ttcttggtgc 2580
agtatatccc tttcgatctt tcttttggtt gcctttacgg ttgagaaatt ggtacttcag 2640
aaatacatat cagaacctgt gagtaattac tattctccag ccattactgt aatttttatt 2700
gaagacaagt ttgtatcatg aagaacttac aagttctgtt ttgaaaatgc tcaaggttgt 2760
catctttctt catattatta tcaccatgac agaggttttg tatccagttt acgtcaccct 2820
aaggtgatac tgtttttctg gtctcagttt gtgatactgt ttttaagttt agttgtctga 2880
cccggtgact ttgaaaatgg acaggtgtga ttctgctttt ttatcaggtg tcactttgat 2940
gctcctcact tgcatttgtt ggctaaaagt ggtttcttat gctcatacta gctatgacat 3000
aagatcccta gccaatgcag ctgataaggt aaaatacgaa aaagaagcgt atgtattagt 3060
cacttgcact gtgttactgt ttttaacaaa cactgttatg aactttaggc caatcctgaa 3120
gtctcctact acgttagctt gaagagcttg gcatatttca tggtcgctcc cacatttgtg 3180

```

tatcaggtaa ctgcaaagt catcaacccat tcttatactt gcaagagttt cttgtctaaa 3240
 cctcggatct ttgcttttcc ccagccaagt tatccacgtt ctgcatgtat acggaagggg 3300
 tgggtggctc gtcaatttgc aaaactgggc atattcacgc gattcatggg atttataata 3360
 gaacaagtac gttttcacat cttgctttat tagttttcct tgggtgaaaat catcatccct 3420
 gcgttgctac cacttgactt catgttcttt tgttacattt tggcagtata taaatcctat 3480
 tgtcaggaac tcaaagcatc ctttgaaagg cgatcttcta tatgctattg aaagagtgtt 3540
 gaagctttca gttccaaatt tatatgtgtg gctctgcatg ttctactgct tcttccacct 3600
 ttggtatgct gtgatcccat ctctttcaaa ataatttgca aattcgaaaa accgaaaaag 3660
 gctaaatctc atacgaattt gatattttta gtttcttaga gtcgggtgat taatttcagt 3720
 tactgaacgc aaatctcttg tccaaagggt aaacatattg gcagagcttc tctgcttcgg 3780
 ggatcgtgaa ttctacaaag attggtggaa tgcaaaaagt gtgggagatg tgagctattt 3840
 tactcaaaag aaaacttatg atttttaatg ttgtcgttgt ttttgggtca tctaactaac 3900
 caaattcatg tattcactgt ctctctttat cagtactgga gaatgtggaa tatggtatgg 3960
 ttctcttctc aaacatcacc ttcttttgta cacaaaatag aagaagagag ctaattaaga 4020
 tcttggtttc cttgacagcc tgttcataaa tggatgggtc gacatatata cttcccgtgc 4080
 ttgcgagca agataccaaa ggtgagtgtg atatataccg atatgcaatt gtcgagattt 4140
 gtttctgtga tataaattta accctccaca cacttgtttt tcagacactc gccattatca 4200
 ttgctttcct agtctctgca gtctttcatg aggtatacat actttctaca ttgccctgtc 4260
 tctagacgca tgaacacacg ctagtgaag aaatgctaatt attcaaagca ttgtttttac 4320
 ttaacgatct tgtgttataa atttcctttt gacagctatg catcgcagtt ccttgtcgtc 4380
 tcttcaagct atgggctttt cttgggatta tgtttcaggt taaaaaatta ctaaactgct 4440
 gcagtcgatt tttactaaac tctaacttca tattctgacc aaccaatttg tttgagtagg 4500
 tgcttttggt cttcatcaca aactatctac aggaaagggt tggctcaacg gtatgctctc 4560
 aaaacccgag aaaatagaac gaataactct ttctttcata gcctagccat ttaaactgca 4620
 atgctgaaac ttaataataa aggtgatctg ttttggaaatg ggatcatatt attaggtggg 4680
 gaacatgatc ttctggttca tcttctgcat tttcggacaa ccgatgtgtg tgcttcttta 4740
 ttaccacgac ctgatgaacc gaaaaggatc gatgtcatga aacaactgtt caaaaaatga 4800
 ctttcttcaa acatctatgg cctcgttgga tctccgttga tgttgtggtg gttctgatgc 4860
 taaaacgaca aatagtgtta taaccattga agaagaaaag aaaattagag ttgttgatc 4920
 tgcaaaaatt ttggtagaga cacgcgaacc cgtttggatt ttgttatggt gtaaagaaat 4980
 ttcaatcaaa aaactgttgt aataatttgt accaaaaaga aatgcttttc tggaaacgag 5040
 gggaaaaata gtagttttgt taggttttac tgtttggacc aaatctagta aaaaactttt 5100
 tgtaataagg aaaaaaaaaa aacaaatgtg ataaatgcat ggggattgta tgaaaccttc 5160
 caataaagtt gattggtggt cccgttttgg gga 5193

<210> 4
 <211> 498
 <212> PRT
 <213> mouse

<400> 4
 Met Gly Asp Arg Gly Gly Ala Gly Ser Ser Arg Arg Arg Arg Thr Gly
 1 5 10 15
 Ser Arg Val Ser Val Gln Gly Gly Ser Gly Pro Lys Val Glu Glu Asp
 20 25 30
 Glu Val Arg Asp Ala Ala Val Ser Pro Asp Leu Gly Ala Gly Gly Asp
 35 40 45
 Ala Pro Ala Pro Ala Pro Ala Pro Ala His Thr Arg Asp Lys Asp Gly
 50 55 60
 Arg Thr Ser Val Gly Asp Gly Tyr Trp Asp Leu Arg Cys His Arg Leu
 65 70 75 80

Gln	Asp	Ser	Leu	Phe	Ser	Ser	Asp	Ser	Gly	Phe	Ser	Asn	Tyr	Arg	Gly	
				85					90					95		
Ile	Leu	Asn	Trp	Cys	Val	Val	Met	Leu	Ile	Leu	Ser	Asn	Ala	Arg	Leu	
				100					105					110		
Phe	Leu	Glu	Asn	Leu	Ile	Lys	Tyr	Gly	Ile	Leu	Val	Asp	Pro	Ile	Gln	
				115					120					125		
Val	Val	Ser	Leu	Phe	Leu	Lys	Asp	Pro	Tyr	Ser	Trp	Pro	Ala	Pro	Cys	
				130					135					140		
Val	Ile	Ile	Ala	Ser	Asn	Ile	Phe	Val	Val	Ala	Ala	Phe	Gln	Ile	Glu	
				145					150					155		
Lys	Arg	Leu	Ala	Val	Gly	Ala	Leu	Thr	Glu	Gln	Met	Gly	Leu	Leu	Leu	
				165					170					175		
His	Val	Val	Asn	Leu	Ala	Thr	Ile	Ile	Cys	Phe	Pro	Ala	Ala	Val	Ala	
				180					185					190		
Leu	Leu	Val	Glu	Ser	Ile	Thr	Pro	Val	Gly	Ser	Val	Phe	Ala	Leu	Ala	
				195					200					205		
Ser	Tyr	Ser	Ile	Met	Phe	Leu	Lys	Leu	Tyr	Ser	Tyr	Arg	Asp	Val	Asn	
				210					215					220		
Leu	Trp	Cys	Arg	Gln	Arg	Arg	Val	Lys	Ala	Lys	Ala	Val	Ser	Thr	Gly	
				225					230					235		
Lys	Lys	Val	Ser	Gly	Ala	Ala	Ala	Gln	Gln	Ala	Val	Ser	Tyr	Pro	Asp	
				245					250					255		
Asn	Leu	Thr	Tyr	Arg	Asp	Leu	Tyr	Tyr	Phe	Ile	Phe	Ala	Pro	Thr	Leu	
				260					265					270		
Cys	Tyr	Glu	Leu	Asn	Phe	Pro	Arg	Ser	Pro	Arg	Ile	Arg	Lys	Arg	Phe	
				275					280					285		
Leu	Leu	Arg	Arg	Val	Leu	Glu	Met	Leu	Phe	Phe	Thr	Gln	Leu	Gln	Val	
				290					295					300		
Gly	Leu	Ile	Gln	Gln	Trp	Met	Val	Pro	Thr	Ile	Gln	Asn	Ser	Met	Lys	
				305					310					315		
Pro	Phe	Lys	Asp	Met	Asp	Tyr	Ser	Arg	Ile	Ile	Glu	Arg	Leu	Leu	Lys	
				325					330					335		
Leu	Ala	Val	Pro	Asn	His	Leu	Ile	Trp	Leu	Ile	Phe	Phe	Tyr	Trp	Phe	
				340					345					350		
Phe	His	Ser	Cys	Leu	Asn	Ala	Val	Ala	Glu	Leu	Leu	Gln	Phe	Gly	Asp	
				355					360					365		
Arg	Glu	Phe	Tyr	Arg	Asp	Trp	Trp	Asn	Ala	Glu	Ser	Val	Thr	Tyr	Phe	
				370					375					380		

435

440

445

Gln Gly Asn Tyr Gly Asn Ala Ala Val Trp Leu Ser Leu Ile Ile Gly
 450 455 460

Gln Pro Ile Ala Val Leu Met Tyr Val His Asp Tyr Tyr Val Leu Asn
 465 470 475 480

Tyr Glu Ala Pro Ala Ala Glu Ala
 485

<210> 6
 <211> 629
 <212> DNA
 <213> Arabidopsis thaliana

<220>
 <221>
 <222>
 <223>

etc ...

SECTIONS
 DEFINING
 MISC FEATURE
 IE "n" in following
 SEQUENCE.

<400> 6
 tgcattgtata cggaagggtt ggggtggctcg tcaatttgca aaactgggtca tattcaccgg 60
 attcatggga tttataatag aacaatatat aaatcctatt gtcaggaact caaagcatcc 120
 tttgaaaggc gatcttctat atgctattga aagagtgttg aagctttcag ttccaaattt 180
 atatgtgtgg ctctgcatgt tctactgctt cttccacctt tggttaaaca tattggcaga 240
 gcttctctgc ttcggggacg gtgaattcta caaagattgg tggaatgcaa aaagtgtggg 300
 agattactgg gagaatgtgg aatatgcctg tccataaatg ggatgggtcc gacatatata 360
 ccttccccgt gcttgccac aaggattacc caaagacacc ccggccatta accattggct 420
 ttccaagcc cctggaggc ctttccatgg gccanggacc cggngtnccc tggcngggcc 480
 ttcaaagcaa agggggnttn cctggggnta aagntccang ggcccttggg gccanccaa 540
 aannttcccc cgggaaaggg ttgccaccg gggggngaaa aanncccggg ggcaccncgg 600
 aattttggga acccgggggg ggccttttt 629

<210> 7
 <211> 11
 <212> PRT
 <213> Arabidopsis thaliana

<400> 7
 His Lys Trp Met Val Arg His Ile Tyr Phe Pro
 1 5 10

<210> 8
 <211> 341
 <212> PRT
 <213> Brassica napus

<400> 8
 Met Cys Cys Leu Ser Leu Ser Ile Phe Pro Leu Ala Ala Phe Thr Val
 1 5 10 15

Glu Lys Leu Val Leu Gln Lys Cys Ile Ser Glu Pro Val Val Ile Phe
 20 25 30

Leu His Val Ile Ile Thr Met Thr Glu Val Leu Tyr Pro Val Tyr Val
 35 40 45

FOUO "4152960

Lys Gly Ser Met Ser
340

<211> 503

<212> PRT

<213> Brassica napus

<400> 9

Met Ala Ile Leu Asp Ser Gly Gly Val Ala Val Pro Pro Thr Glu Asn
1 5 10 15

Gly Val Ala Asp Leu Asp Arg Leu His Arg Arg Lys Ser Ser Ser Asp
20 25 30

Ser Ser Asn Gly Leu Leu Ser Asp Thr Ser Pro Ser Asp Asp Val Gly
35 40 45

Ala Ala Ala Ala Glu Arg Asp Arg Val Asp Ser Ala Ala Glu Glu Glu
50 55 60

Ala Gln Gly Thr Ala Asn Leu Ala Gly Gly Asp Ala Glu Thr Arg Glu
65 70 75 80

Ser Ala Gly Gly Asp Val Arg Phe Thr Tyr Arg Pro Ser Val Pro Ala
85 90 95

His Arg Arg Thr Arg Glu Ser Pro Leu Ser Ser Asp Ala Ile Phe Lys
100 105 110

Gln Ser His Ala Gly Leu Phe Asn Leu Cys Val Val Val Leu Val Ala
115 120 125

Val Asn Ser Arg Leu Ile Ile Glu Asn Leu Met Lys Tyr Gly Trp Leu
130 135 140

Ile Arg Thr Asp Phe Trp Phe Ser Ser Thr Ser Leu Arg Asp Trp Pro
145 150 155 160

Leu Phe Met Cys Cys Leu Ser Leu Ser Val Phe Pro Leu Ala Ala Phe
165 170 175

Thr Val Glu Lys Met Val Leu Gln Lys Phe Ile Ser Glu Pro Val Ala
180 185 190

Ile Ile Leu His Val Ile Ile Thr Met Thr Glu Val Leu Tyr Pro Val
195 200 205

Tyr Val Thr Leu Arg Cys Asp Ser Ala Phe Leu Ser Gly Val Thr Leu
210 215 220

Met Leu Leu Thr Cys Ile Val Trp Leu Lys Leu Val Ser Tyr Ala His
225 230 235 240

Thr Ser Tyr Asp Ile Arg Thr Leu Ala Asn Ser Ala Asp Lys Val Asp
245 250 255

Pro Glu Ile Ser Tyr Tyr Val Ser Leu Lys Ser Leu Ala Tyr Phe Met
 260 265 270

Val Ala Pro Thr Leu Cys Tyr Gln Pro Ser Tyr Pro Arg Ser Pro Cys
 275 280 285

Ile Arg Lys Gly Trp Val Ala Arg Gln Leu Ala Lys Leu Val Ile Phe
 290 295 300

Thr Gly Leu Met Gly Phe Ile Ile Glu Gln Tyr Ile Asn Pro Ile Val
 305 310 315 320

Arg Asn Ser Lys His Pro Leu Lys Gly Asp Leu Leu Tyr Ala Ile Glu
 325 330 335

Arg Val Leu Lys Leu Ser Val Pro Asn Leu Tyr Val Trp Leu Cys Met
 340 345 350

Phe Tyr Cys Phe Phe His Leu Trp Leu Asn Ile Leu Ala Glu Leu Leu
 355 360 365

Cys Phe Gly Asp Arg Glu Phe Tyr Lys Asp Trp Trp Asn Ala Lys Ser
 370 375 380

Val Gly Asp Tyr Trp Arg Met Trp Asn Met Pro Val His Lys Trp Met
 385 390 395 400

Val Arg His Val Tyr Phe Pro Cys Leu Arg Ile Lys Ile Pro Lys Val
 405 410 415

Pro Ala Ile Ile Ile Ala Phe Leu Val Ser Ala Val Phe His Glu Leu
 420 425 430

Cys Ile Ala Val Pro Cys Arg Leu Phe Asn Leu Trp Ala Phe Met Gly
 435 440 445

Ile Met Phe Gln Val Pro Leu Val Phe Ile Thr Asn Phe Leu Gln Glu
 450 455 460

Arg Phe Gly Ser Met Val Gly Asn Met Ile Phe Gly Ser Ala Ser Cys
 465 470 475 480

Ile Phe Gly Gln Pro Met Cys Gly Leu Leu Tyr Tyr His Asp Leu Met
 485 490 495

Asn Arg Lys Gly Ser Met Ser
 500

<210> 10
 <211> 11
 <212> PRT
 <213> Arabidopsis thaliana

[<220>
 <221> VARIANT]

<220>
 <221>
 <222>
 <223>
 etc..

SECTIONS
 DEFINING
 MISC FEATURES
 I.E. "Xaa" in
 FOLLOWING
 SEQUENCE.

<222> (1)

<220>

<221> VARIANT

<222> (3)

<220>

<221> VARIANT

<222> (5)..(6)

<220>

<221> VARIANT

<222> (8)..(10)

<400> 10

Xaa Leu Xaa Lys Xaa Xaa Ser Xaa Xaa Xaa Val
1 5 10

<210> 11

<211> 27

<212> PRT

<213> Arabidopsis thaliana

<400> 11

Ser His Ala Gly Leu Phe Asn Leu Cys Val Val Val Leu Ile Ala Val
1 5 10 15

Asn Ser Arg Leu Ile Ile Glu Asn Leu Met Lys
20 25

<210> 12

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer of
DGATXbal

<400> 12

ctagtctaga atggcgattt tgga

24

<210> 13

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer of
DGATXhol

<400> 13

1062504452950

23

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer DGAT1

<400> 14

agacacgaat cccattccca ccga

24

<210> 15

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer DGAT2

<400> 15

agtgggtgaca acgcaggggat gatg

24

<210> 16

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer DGAT3

<400> 16

atggtcgctc ccacattgtg t

21

<210> 17

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer DGAT4

<400> 17

catacaatcc ccatgacatt tatca

25

<210> 18

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

24

<213> Artificial Sequence

24

<213> Artificial Sequence

21

<213> Artificial Sequence

33

<213> Artificial Sequence

33

<212> DNA

<213> Arabidopsis thaliana

<400> 23

atttcttagc ttcttctctc aatccgctct ttcctctctc attagattct gtttctctct 60
 tcaattttct ctgcattgct ctgcattctc tctgacgcct cttttctccc gacgctgttt 120
 cgtcaaacgc ttttcgaaat ggcgattttg gattctgctg gcgttactac ggtgacggag 180
 aacggtggcg gagagtctgt cgatcttgat aggcctcgtc gacggaaatc gagatcggat 240
 tcttctaacg gacttctctc ctctggttcc gataataatt ctcttcgga tgatgttga 300
 gctcccgccg acgttaggga tccgattgat tccgttgta acgatgacgc tcagggaaca 360
 gccaatgtg ccggagataa taacggtggt ggcgataata acggtggtg aagaggcgcc 420
 ggagaaggaa gaggaacgc cgatgctacg tttacgtatc gacgctcgtt tccagctcat 480
 cggaggcgga gagagagtc acttagctcc gacgcaatct tcaaacagag ccatgccgga 540
 ttattcaacc tctgtgtagt agttcttatt gctgtaaaca gtagactcat catcgaaaat 600
 cttatgaaga gccatgccgg attattcaac ctctgtgtag tagttcttat tgctgtaaac 660
 agtagactca tcatcgaaaa tcttatgaag tatggttggt tgatcagaac ggatttctgg 720
 tttagttcaa gatcgctgcg agattggccg cttttcatgt gttgtatata ctttctgac 780
 tttcttttgg ctgcttttac ggttgagaaa ttggtacttc agaaatacat atcagaacct 840
 gttgtcatct ttcttcatat tattatcacc atgacagagg ttttgtatcc agtttacgtc 900
 accctaaggt gtgattctgc ttttttatca ggtgtcact tgatgctcct cacttgcat 960
 gtgtggctaa agttggtttc ttatgctcat actgctatg acataagatc ctagccaat 1020
 gcagctgata aggccaatcc tgaagtctcc tactacgtta gcttgaagag cttggcatat 1080
 ttcatggtcg ctcccacatt gtgttatcag ccaagttatc cacgttctgc atgtatacgg 1140
 aagggttggg tggctcgtca atttgcaaaa ctggtcatat tcaccggatt catgggattt 1200
 ataatagaac aatatataaa tctattgtc aggaactcaa agcatccttt gaaaggcgat 1260
 cttctatatg ctattgaaag agtgttgaag ctttcagttc caaatattata tgtgtggctc 1320
 tgcattgtct actgcttctt ccacctttgg ttaaacatat tggcagagct tctctgcttc 1380
 ggggatcgtg aattctacaa agattgggtg aatgcaaaaa gtgtgggaga ttactggaga 1440
 atgtggaata tgctgttca taaatggatg gttcgacata tatacttccc gtgcttgcgc 1500
 agcaagatac caaagacact cgccattatc attgctttcc tagtctctgc agtctttcat 1560
 gagctatgca tgcagttcc ttgtcgtctc ttcaagctat gggcttttct tgggattatg 1620
 tttcaggtgc ctttggctct catcacaac tatctacagg aaagggttgg ctcaacggtg 1680
 gggaaacatga tcttctggtt catcttctgc attttcggac aaccgatgtg tgtgcttctt 1740
 tattaccacg acctgatgaa ccgaaaagga tccatgtcat gaaacaactg ttcaaaaaat 1800
 gactttcttc aaacatctat ggctcgttg gatctccgtt gatgttgtgg tggttctgat 1860
 gctaaaacga caaatagtgt tataaccatt gaagaagaaa agaaaattag agttgttgta 1920
 tctgcaaaaa ttttggtaga gacacgcaaa cccgtttgga ttttgttatg gtgtaaagcg 1980
 gccgc 1985

<210> 24

<211> 5339

<212> DNA

<213> Arabidopsis thaliana

<400> 24

gctcacgacc cattcttccc gttccatttg gttttattta tttcaaagtt taatattcct 60
 tttgtataac attcaaatct tcacatgatt gattgtgtga aaacccaca gattttacta 120
 caataggggg agttgactta aaatagctat tgatgtcgaa aaaatgtatt ttagttataa 180
 attatactaa agaaaatttt tgatttgtct gttgtttaag catatgtatt gttaaactta 240
 aaaaaatatg tattgttaat cttaaaaatg taggagtaca catcaaatat tcgagcataa 300
 tcaaaaccgt attcatagac cgatgtgaga atcaaataga agataatgtg atttttttaa 360
 atatcgtatc tccaaatcaa tcaattagaa gataatgtaa ttctttatgt gctacataaa 420
 taaatatata tatatatata tatatatatc ttgtatatat gtcttgacaa aaaattgcc 480
 gtcaaaaacc atgactgaat caaactataa gtcggattga atcaaaactat aagtcggatg 540
 agtattaatt tccattatgt ttctatactt tacaaaaccg aaaatagata ttatagatac 600
 caaaaaagta gatttgtgta tattattaga agatttggaa tttcatcatt atcaggatct 660
 aaagtacttc cctaattaaa tcatgtcggg tgaaaaagct caatgaatgt ttgaaatttg 720


```
<210> 25
<211> 547
<212> PRT
<213> Arabidopsis thaliana
```

Met Ala Ile Leu Asp Ser Ala Gly Val Thr Thr Val Thr Glu Asn Gly
1 5 10 15

Gly Gly Glu Phe Val Asp Leu Asp Arg Leu Arg Arg Arg Lys Ser Arg
20 25 30

Ser Asp Ser Ser Asn Gly Leu Leu Leu Ser Gly Ser Asp Asn Asn Ser
35 40 45

Pro Ser Asp Asp Val Gly Ala Pro Ala Asp Val Arg Asp Arg Ile Asp
50 55 60

Ser Val Val Asn Asp Asp Ala Gln Gly Thr Ala Asn Leu Ala Gly Asp
65 70 75 80

Asn Asn Gly Gly Gly Asp Asn Asn Gly Gly Gly Arg Gly Gly Gly Glu
85 90 95

Gly Arg Gly Asn Ala Asp Ala Thr Phe Thr Tyr Arg Pro Ser Val Pro
100 105 110

Ala His Arg Arg Ala Arg Glu Ser Pro Leu Ser Ser Asp Ala Ile Phe
115 120 125

Lys Gln Ser His Ala Gly Leu Phe Asn Leu Cys Val Val Val Leu Ile
130 135 140

Ala Val Asn Ser Arg Leu Ile Ile Glu Asn Leu Met Lys Ser His Ala
145 150 155 160

Gly	Leu	Phe	Asn	Leu	Cys	Val	Val	Val	Leu	Ile	Ala	Val	Asn	Ser	Arg	
				165					170						175	
Leu	Ile	Ile	Glu	Asn	Leu	Met	Lys	Tyr	Gly	Trp	Leu	Ile	Arg	Thr	Asp	
			180					185					190			
Phe	Trp	Phe	Ser	Ser	Arg	Ser	Leu	Arg	Asp	Trp	Pro	Leu	Phe	Met	Cys	
		195					200					205				
Cys	Ile	Ser	Leu	Ser	Ile	Phe	Pro	Leu	Ala	Ala	Phe	Thr	Val	Glu	Lys	
	210					215					220					
Leu	Val	Leu	Gln	Lys	Tyr	Ile	Ser	Glu	Pro	Val	Val	Ile	Phe	Leu	His	
225					230					235					240	
Ile	Ile	Ile	Thr	Met	Thr	Glu	Val	Leu	Tyr	Pro	Val	Tyr	Val	Thr	Leu	
				245					250					255		
Arg	Cys	Asp	Ser	Ala	Phe	Leu	Ser	Gly	Val	Thr	Leu	Met	Leu	Leu	Thr	
			260					265					270			
Cys	Ile	Val	Trp	Leu	Lys	Leu	Val	Ser	Tyr	Ala	His	Thr	Ser	Tyr	Asp	
		275					280					285				
Ile	Arg	Ser	Leu	Ala	Asn	Ala	Ala	Asp	Lys	Ala	Asn	Pro	Glu	Val	Ser	
	290					295					300					
Tyr	Tyr	Val	Ser	Leu	Lys	Ser	Leu	Ala	Tyr	Phe	Met	Val	Ala	Pro	Thr	
305					310					315					320	
Leu	Cys	Tyr	Gln	Pro	Ser	Tyr	Pro	Arg	Ser	Ala	Cys	Ile	Arg	Lys	Gly	
				325					330					335		
Trp	Val	Ala	Arg	Gln	Phe	Ala	Lys	Leu	Val	Ile	Phe	Thr	Gly	Phe	Met	
			340					345					350			
Gly	Phe	Ile	Ile	Glu	Gln	Tyr	Ile	Asn	Pro	Ile	Val	Arg	Asn	Ser	Lys	
		355					360					365				
His	Pro	Leu	Lys	Gly	Asp	Leu	Leu	Tyr	Ala	Ile	Glu	Arg	Val	Leu	Lys	
	370					375					380					
Leu	Ser	Val	Pro	Asn	Leu	Tyr	Val	Trp	Leu	Cys	Met	Phe	Tyr	Cys	Phe	
385					390					395					400	
Phe	His	Leu	Trp	Leu	Asn	Ile	Leu	Ala	Glu	Leu	Leu	Cys	Phe	Gly	Asp	
				405					410					415		
Arg	Glu	Phe	Tyr	Lys	Asp	Trp	Trp	Asn	Ala	Lys	Ser	Val	Gly	Asp	Tyr	
			420					425					430			
Trp	Arg	Met	Trp	Asn	Met	Pro	Val	His	Lys	Trp	Met	Val	Arg	His	Ile	
		435					440					445				
Tyr	Phe	Pro	Cys	Leu	Arg	Ser	Lys	Ile	Pro	Lys	Thr	Leu	Ala	Ile	Ile	
	450					455					460					

Ile Ala Phe Leu Val Ser Ala Val Phe His Glu Leu Cys Ile Ala Val
 465 470 475 480

Pro Cys Arg Leu Phe Lys Leu Trp Ala Phe Leu Gly Ile Met Phe Gln
 485 490 495

Val Pro Leu Val Phe Ile Thr Asn Tyr Leu Gln Glu Arg Phe Gly Ser
 500 505 510

Thr Val Gly Asn Met Ile Phe Trp Phe Ile Phe Cys Ile Phe Gly Gln
 515 520 525

Pro Met Cys Val Leu Leu Tyr Tyr His Asp Leu Met Asn Arg Lys Gly
 530 535 540

Ser Met Ser
 545

050354-03001